



## Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS  
KABI

WATCH VIDEO

# AJNR

## Rapid development of basal ganglia calcification.

M Sarwar and K Ford

*AJNR Am J Neuroradiol* 1981, 2 (1) 103-104

<http://www.ajnr.org/content/2/1/103.citation>

This information is current as  
of August 23, 2025.

# Rapid Development of Basal Ganglia Calcification

Mohammad Sarwar<sup>1, 2</sup> and Kerry Ford<sup>1</sup>

The pathologic and radiologic appearance of calcification of the corpus striatum is now well recognized. This calcification can be associated with endocrine, congenital/developmental, inflammatory, and anoxic/toxic conditions. Although the association of the basal ganglia calcification with these conditions is well established, its pathogenesis is poorly understood. We found no specific answer in the literature as to how long it takes for the pathologic basal ganglia calcification to occur. However, we got the impression that it takes at least several months to years. We recently encountered a young patient in whom the basal ganglia calcification was demonstrated on sequential computed tomography (CT) scans after only 31 days.

## Case Report

A 29-year-old woman was admitted after an alcoholic binge of 2 days. She was nauseated and complained of epigastric pain. Laboratory studies revealed decreased serum levels of sodium, calcium, and phosphate. Her condition deteriorated and she was believed to have hemorrhagic necrotic pancreatitis. An exploratory laparotomy confirming this was complicated by a stormy postoperative course. She developed diabetes mellitus, respiratory difficulty, and acute tubular necrosis.

Serial serum calcium and phosphate levels remained quite low for 2–4 weeks after the operation (serum calcium levels were as low as 4–5 mg/dl and serum phosphate levels as low as 1 mg/dl). During this time she sustained two cardiopulmonary arrests with successful resuscitation. Postoperative neurologic status varied from decorticate and decerebrate posturing to the ability to speak only short sentences. Whereas she could execute spontaneous movements in all her extremities, her tone was considerably increased and she had tremors of both extremities (extrapyramidal signs). Her other medical problems were adequately controlled.

An initial CT scan at admission was normal (fig. 1A). A scan 31 days later showed extensive basal ganglia calcification (fig. 1B). A lateral skull film done 3 weeks after the second scan revealed no basal ganglia calcification.

## Discussion

Basal ganglia calcification was initially reported independently by Virchow [1] and Bamberger [2] in 1855. Its radio-

graphic visualization was first described by Fritzsche [3] in 1935. Since the reports of Eaton and Haines [4] and Sprague et al. [5], who showed relation of basal ganglia calcification to hypoparathyroidism and pseudohypoparathyroidism, respectively, many other reports have discussed the association of basal ganglia calcifications with other conditions [6–17]. From these reports it is clearly evident that in the central nervous system, the basal ganglia, particularly the globus pallidus, show a great proclivity for calcification. Our case further supports this premise. From the clinical summary it is perhaps fair to conclude that the underlying cause of basal ganglia calcification in our case was anoxia.

The vulnerability of the basal ganglia, particularly of the globus pallidus, to anoxic injury (pallidal necrosis) has long been recognized [17, 18]. It is rare for the corpus striatum (putamen and caudate nucleus) to suffer this type of anoxic injury without concomitant pallidal injury. The *pathoclasia* (susceptibility to involvement by toxins) theory of Vogt and Vogt [19] was based on the assumption that certain parts of the brain had physiochemical properties that made them more susceptible to anoxic damage [20, 21]. Several factors have been postulated to explain this susceptibility of the globus pallidus and corpus striatum to anoxic injury and consequent dystrophic calcification. These factors include:




Fig. 1.—A, Normal admission scan. B, 31 days later. Extensive basal ganglia calcification.

Received June 24, 1980; accepted after revision August 19, 1980

<sup>1</sup>Department of Radiology, University of Texas Medical Branch at Galveston, Galveston, TX 77550.

<sup>2</sup>Present address: Department of Radiology, Yale University School of Medicine, 333 Cedar St., New Haven, CT 06510. Address reprint requests to M. Sarwar.

AJNR 2:103–104, January/February 1981 0195–6108/81/0021–0103 \$00.00 © American Roentgen Ray Society



(1) abundance of oxidative enzymes [20–22]; (2) peculiarities of the arterial supply [23–29]; and (3) disturbance of vasomotor regulation [30]. Physiologic calcification of the globus pallidus after the age of 40–50 years gives further credence to this theory [16]. It is difficult to ascertain if the relative abundance of iron in the globus pallidus increases its susceptibility to calcium deposition [31].

Pathologically the calcification is located in and around the finer blood vessels [32]. This calcification follows colloid deposition in the substrate; the colloid then changes into a gel that accepts calcium salts [32].

We were unable to find information regarding the time required for secondary basal ganglia calcification to occur. However, our case suggests that the basal ganglia are not only highly susceptible to anoxic injury and consequent dystrophic calcification, but also that basal ganglia calcification can occur in as short a time as 31 days, much more rapidly than generally believed.

#### ACKNOWLEDGMENT

We thank Ramona Guzman for manuscript preparation.

#### REFERENCES

- Virchow R. Kalk-Metastases. *Virchows Arch [Pathol Anat]* **1855**;8:103–133
- Bamberger PH. Beobachtungen und Bemerkungen über Hirnkrankheiten. *Verh Phys Med Ges Würzburg* **1855**;6:325–328
- Fritzche R. Eine familiär auftretende Form von oligophrenie mit röntgenologisch nachweisbaren symmetrischen Kalkablagerungen in Gehirn, besonders in den Stammganglien. *Schweiz Arch Neurol Neurochir Psychiatr* **1935**;35:1–29
- Eaton LM, Haines SF. Parathyroid insufficiency with symmetrical cerebral calcification. *JAMA* **1939**;113:749–753
- Sprague RG, Haines SF, Power MH. Metabolic effects of parathyroid hormone, dihydrotachysterol and calciferol in case of pseudohypoparathyroidism. *Pro Centr Soc Clin Res* **1944**;17:16–17
- Camp JD. Symmetrical calcification of cerebral basal ganglia; its roentgenologic significance in diagnosis of parathyroid insufficiency. *Radiology* **1947**;49:568–577
- Camp JD. Pathologic non-neoplastic intracranial calcification. *JAMA* **1948**;137:1023–1031
- Wagner JA, Slager UT, Dennis JM, et al. The incidence and composition of radiopaque deposit in the basal ganglia of the brain. *AJR* **1955**;74:232–234
- Bennett JC, Maffly RH, Steinback HL. The significance of bilateral basal ganglia calcification. *Radiology* **1959**;72:368–378
- Palubinskas AJ, Davies H. Calcification of the basal ganglia of the brain. *AJR* **1959**;82:806–822
- Steinback HL, Young DA. The roentgen appearance of pseudohypoparathyroidism (PH) and pseudo-pseudohypoparathyroidism (PPH). Differentiation from other syndromes associated with short metacarpals, metatarsals, and phalanges. *AJR* **1966**;97:49–66
- Babbitt DP, Tang T, Dobbs J, Berk R. Idiopathic familial cerebrovascular ferrocalsinosis (Fahr's disease) and review of differential diagnosis of intracranial calcification in children. *AJR* **1969**;105:352–358
- Harwood-Nash DC, Reilly BJ. Calcification of the basal ganglia following radiation therapy. *AJR* **1970**;108:392–395
- Lee KF, Suh JH. CT evidence of grey matter calcification secondary to radiation therapy. *Comput Tomogr* **1977**;1:103–110
- Seigel RS, Seeger JF, Gabrielsen TO, et al. Computed tomography in oculocraniosomatic disease (Kearns-Sayre syndrome). *Radiology* **1979**;130:159–164
- Cohen CR, Duchesneau PM, Weinstein MA. Calcification of the basal ganglia as visualized by computed tomography. *Radiology* **1980**;134:97–99
- Lowenthal A, Bruyn GW. Calcification of the striopallidodentate system. In: Vinken PJ, Bruyn GW, eds. *Handbook of clinical neurology*, vol 6. Amsterdam: North-Holland, **1968**:703–729
- Kellinger K. Degenerations and exogenous lesions of the pallidum and striatum. In: Vinken PJ, Bruyn GW, eds. *Handbook of Clinical Neurology*, vol 6. Amsterdam: North-Holland, **1968**:652–654
- Vogt C, Vogt O. Erkrankungen der Grobhirnrinde im Lichte der Topistik Pathoklise und Pathoarchitektonik. *Psychol Neurol* **1922**;28:1–71
- Friede RL. Chemoarchitecture and neuropathology. In: *Proc IVth Int. Congr. Neuropath*, vol 20. Stuttgart: Thieme, **1962**:70–75
- Jacob H. Pattern of CNS-vulnerability-CNS tissue and cellular pathology states. In: Schage JP, McMenemey WH, eds. *Selective vulnerability of the brain in hypoxaemia*. Oxford: Blackwell, **1963**:153–163
- Shimizu N, Morikawa N. Histochemical studies of succinic dehydrogenase of the brain of mice, rats, guinea pigs and rabbits. *J Histochem Cytochem* **1957**;5:334–345
- Hiller F. Über die krankhaften Veränderungen im Zentralnervensystem nach Kohlenoxydvergiftung. *Z Ges Neurol Psychiatr* **1924**;93:594–646
- Spielmeyer W. Über örtliche Vulnerabilität. *Z Ges Neurol Psychiatr* **1928**;118:1–16
- Lindenberg R. Compression of brain arteries as pathogenic factor for tissue necroses and their areas of predilection. *J Neuropathol Exp Neurol* **1955**;14:223–243
- Lindenberg R. Patterns of CNS vulnerability in acute hypoxaemia, including anaesthesia accidents. In: Schade JP, McMenemey WH, eds. *Selective vulnerability of the brain in hypoxaemia*. Oxford: Blackwell, **1963**:189–209
- Norman RM, Ulrich H. The influence of a vascular factor on the distribution of symmetrical cerebral calcifications. *J Neurol Neurosurg Psychiatry* **1960**;23:142–147
- Kornyey S. Patterns of CNS vulnerability in CO, cyanide and other poisonings. In: Schade JP, McMenemey WH, eds. *Selective vulnerability of the brain in hypoxaemia*. Oxford: Blackwell, **1963**:165–181
- Alexander L. The vascular supply of the striopallidum. *Res Publ Assoc Nerv Ment Dis* **1942**;21:77–132
- Scholz W. Selective neuronal necrosis and its topistic patterns in Hypoxemia and oligemia. *J Neuropathol Exp Neurol* **1953**;12:249–261
- Neumann MA. Iron and calcium dysmetabolism in the brain. With special predilection for globus pallidus and cerebellum. *J Neuropathol Exp Neurol* **1963**;22:148–163
- Eaton LM, Camp JD, Love JC. Symmetric cerebral calcification, particularly of the basal ganglia, demonstrable roentgenographically. Calcification of the finer blood vessels. *Arch Neurol Psychiatr* **1939**;41:921–942