



Providing Choice & Value
Generic CT and MRI Contrast Agents



CONTACT REP

AJNR

Reply:

Q. Wang, X. Xu and M. Zhang

AJNR Am J Neuroradiol 2010, 31 (9) E81

doi: <https://doi.org/10.3174/ajnr.A2188>

<http://www.ajnr.org/content/31/9/E81>

This information is current as
of July 18, 2025.

Reply:

We thank Dr Hasan for his comments on our study.¹

Brain development and aging are complex processes. During brain maturation, the mean diffusivity (MD) decreases and the fractional anisotropy (FA) increases with age.²⁻⁴ Decreases of MD and increases of FA occur rapidly at young ages, then slow and gradually reach a plateau.³ In most structures, including the genu and splenium of the corpus callosum, the basal ganglia, and so forth, the plateau is reached during the late teens or twenties.³ From young-to-old adults, the diffusivity in the brain presents in a different way from brain maturation. MD increases and FA decreases with age in some structures of the brain.⁵⁻⁷ Thus the linear regression is obviously not appropriate for studying the diffusivity of the brain across the whole lifespan. In our study, all volunteers were adults. We did not include children. For studying age-related alterations in the human brain of young and old adults with diffusion tensor imaging, the Pearson correlation analysis may be simple, but it is acceptable. The trend toward diffusion tensor imaging metric changes with age is not altered just because of the simple model used.⁶⁻⁸

We noticed that the finding of decreased MD with age in the putamen was contradictory to several previous reports.⁵⁻⁷ However, in our study, the results of MD in the prefrontal white matter and the genu of the corpus callosum are consistent with the above-mentioned reports. Many factors may relate to this kind of discrepancy, such as different imaging protocols, imaging planes, the region of interest, and, most of all, the exclusion criteria of the healthy volunteers.

In this article,¹ we did not state that the diffusion tensor eigenvalues can be used as surrogate neuroimaging markers of human brain aging. The information from axial (λ_1) and radial (λ_{23}) diffusivity was the direct contributor to the FA. The availability of this information would yield insight into the microstructural changes of human brain aging.

Acknowledgments

M. Zhang was supported by National Science Foundation of China (No. 30570536). X. Xu was supported by National Science Foundation of China (No. 30900358) and the Medical Scientific Research Foundation of Zhejiang Province, China (No. 2009QN005).

References

1. Wang Q, Xu X, Zhang M. Normal aging in the basal ganglia evaluated by eigenvalues of diffusion tensor imaging. *AJNR Am J Neuroradiol* 2010;31:516-20
2. Snook L, Paulson LA, Roy D, et al. Diffusion tensor imaging of neurodevelopment in children and young adults. *Neuroimage* 2005;26:1164-73
3. Lebel C, Walker L, Leemans A, et al. Microstructural maturation of the human brain from childhood to adulthood. *Neuroimage* 2008;40:1044-55
4. Mukherjee P, Miller JH, Shimony JS, et al. Diffusion-tensor MR imaging of gray and white matter development during normal human brain maturation. *AJNR Am J Neuroradiol* 2002;23:1445-56
5. Pfefferbaum A, Adalsteinsson E, Rohlfing T, et al. Diffusion tensor imaging of deep gray matter brain structures: effects of age and iron concentration. *Neurobiol Aging* 2010;31:482-93
6. Salat DH, Tuch DS, Greve DN, et al. Age-related alterations in white matter microstructure measured by diffusion tensor imaging. *Neurobiol Aging* 2005;26:1215-27
7. Abe O, Aoki S, Hayashi N, et al. Normal aging in the central nervous system: quantitative MR diffusion-tensor analysis. *Neurobiol Aging* 2002;23:433-41
8. Abe O, Yamasue H, Aoki S, et al. Aging in the CNS: comparison of gray/white matter volume and diffusion tensor data. *Neurobiol Aging* 2008;29:102-16

Q. Wang

X. Xu

M. Zhang

Department of Radiology

First Affiliated Hospital

School of Medicine, Zhejiang University

Hangzhou, China

DOI 10.3174/ajnr.A2188