



Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Dementia, Quantitative Neuroimaging, and Apolipoprotein E Genotype

Erin D. Bigler, Christopher M. Lowry, Carol V. Anderson, Sterling C. Johnson, John Terry and Marc Steed

AJNR Am J Neuroradiol 2000, 21 (10) 1857-1868

<http://www.ajnr.org/content/21/10/1857>

This information is current as
of August 8, 2025.

Dementia, Quantitative Neuroimaging, and Apolipoprotein E Genotype

Erin D. Bigler, Christopher M. Lowry, Carol V. Anderson, Sterling C. Johnson, John Terry, and Marc Steed

BACKGROUND AND PURPOSE: Quantitative MR imaging differences in an elderly population of subjects with various clinical disorders (including dementia, particularly Alzheimer's disease and vascular dementia) and disorders of mild cognitive impairment were examined. Potential quantitative MR differences were assessed by presence or absence of the apolipoprotein E (*APOE*) $\epsilon 4$ allele and by level of cognitive deficit.

METHODS: One hundred eighty subjects with a diagnosis of dementia or other clinical disorders were identified from an eligible population of 5677 elderly individuals. Age, duration of disease, and head size (where appropriate) were considered as covariates. *APOE* genotype was determined by polymerase chain reaction using buccal material. Axial and coronal intermediate- and T2-weighted MR images were quantified using a multispectral segmentation algorithm. Cognitive status was assessed by means of a modified Mini-Mental Status Examination.

RESULTS: All types of dementing illness showed significant volume reductions in the majority of structures examined, particularly in the total brain, hippocampus, and white and gray matter, and increased CSF and ventricular volumes. Subjects with mild cognitive impairment showed fewer atrophic changes but were still distinguishable from the 24 control subjects. Presence of an $\epsilon 4$ allele was associated with smaller hippocampal volume in subjects with Alzheimer's disease and vascular dementia within just 1 year of disease onset. For other analyses, atrophy related to the presence of the $\epsilon 4$ allele disappeared after controlling for age and length of disease.

CONCLUSION: The effects of the $\epsilon 4$ allele on brain morphology may be subtly expressed early in the development of dementia, but do not specifically affect cerebral atrophy thereafter. Cognitive impairment is associated with atrophy irrespective of diagnosis and presence of $\epsilon 4$.

A major public health concern is aging of the population and associated increases in the prevalence of various dementias, particularly Alzheimer's disease (AD) (1–3). Neuroimaging is a key procedure in the assessment of dementia, especially for differential diagnostic evaluations (4–10). Recently, interest has focused on quantitative methods for distinguishing normal aging from pathologic conditions and for differentiating the types of dementia, such as AD versus vascular dementia (VaD) (5, 6, 10–17). Few investigators, however, have had the opportunity to examine an entire population.

Recent dementia research has focused on the genetics of degenerative diseases (18). For example, numerous studies have shown an increased risk for

AD, and potentially for VaD, associated with the allele $\epsilon 4$ at *APOE*, the polymorphic genetic locus for apolipoprotein E (19–33), but the relationship between AD, $\epsilon 4$, and neuroimaging findings remains uncertain (1, 2, 5, 10).

If the $\epsilon 4$ allele is associated with increased risk for AD and VaD, does it also predict degenerative, atrophic changes that are detectable by quantitative MR imaging techniques? To date, results in this area have been equivocal. In a preliminary study that included only a subset of the AD subjects reported herein, we found the $\epsilon 4$ allele to be associated with smaller total brain and hippocampal volumes, as well as with a larger ventricle-to-brain ratio (VBR) (34). Additionally, diminished cognitive performance (eg, on the modified Mini-Mental Status Examination, or 3MS) was related to degree of atrophy. However, when age and length of disease (LOD) were included as covariates, these $\epsilon 4$ -associated differences in brain morphology were no longer evident. Jack et al (5, 35) recently reported no apparent *APOE* effect on hippocampal or temporal lobe volume, and Yasuda et al (20) found that

Received January 3, 2000; accepted after revision May 23.

From the Department of Psychology and Neuroscience, 1001 SWKT, Brigham Young University, Provo, UT 84602. Address reprint requests to Erin D. Bigler, PhD.

Supported in part by NIH grants AG-11380 and MH-14592, and by the Ira Fulton Foundation.

AD patients with homozygous $\epsilon 4$ had the least amount of atrophy. Barber et al (36, 37) did not find medial temporal lobe atrophy or white matter lesions to be associated with *APOE* genotype. In contrast, Lehtovirta et al (38, 39) and Soininen and coworkers (40, 41) found smaller hippocampal volumes in AD subjects with an $\epsilon 4$ allele and smaller temporal lobe and associated structures that were $\epsilon 4$ dose-dependent (42). Plassman et al (43) also found smaller hippocampal volumes in nondemented, cognitively intact subjects with the $\epsilon 4$ allele, consistent with observations by Reiman et al (8) and Tohgi et al (44). None of these studies was population based, so sample variations may explain some of the discrepancy in the findings. Also, cerebral blood flow (CBF) studies have shown conflicting relationships between CBF and *APOE* genotype (45–49).

A new epidemiologic investigation of aging and dementia in the residents of Cache County, Utah, provided an opportunity to examine quantitative MR findings in an elderly population (including centenarians) with a continuum of cognitive functioning, all of whom had established *APOE* genotypes (50, 51). From a population of 5677 eligible elderly residents in the county, 5092 (90%) underwent a multistage evaluation process to detect and identify prevalent cases of AD, VaD, and other dementias (29) as well as cases of mild cognitive impairment (MCI) (52). The study also identified subjects with other clinical disorders that may affect cognition, such as Parkinson's disease, stroke, and neuropsychiatric disorders. A particular objective was to investigate the relationship between the *APOE* genotype and dementia. The increased AD risk associated with the $\epsilon 4$ allele was confirmed in the Cache County sample, although the association appeared to wane in extreme old age (50). A trend was also observed between the $\epsilon 4$ allele and VaD.

When possible, the dementia workup in the Cache County study included MR imaging of the brain. The MR images were interpreted clinically and then subjected to quantitative analysis (11, 12). We report herein the quantitative MR analyses in the first 180 patients with AD, VaD, and other clinical disorders, as well as in the 24 cognitively normal subjects, all with known *APOE* genotypes (53). The objectives of this investigation were fourfold: 1) to describe the quantitative findings of MR analyses in this population of subjects with various types of dementing illnesses and known *APOE* genotype; 2) to determine whether the $\epsilon 4$ allele was associated with any unique differences in brain morphology in this population of elderly and demented subjects; 3) to examine age and LOD variables on brain morphology in terms of the presence or absence of the $\epsilon 4$ allele and diagnostic classification; and 4) to examine quantitative MR findings, *APOE* classification, and cognitive impairment as assessed by a modified 3MS examination (54).

Methods

The Cache County Study

A detailed description of the Cache County elderly population and the study methods used have been published elsewhere (50, 51). Briefly, a multistage screening and assessment protocol to ascertain the population's prevalence of dementia was used. This detection method is believed to yield 85% to 90% sensitivity (55). Individuals with dementia underwent a differential diagnostic evaluation that included a detailed history, a brief physical assessment, a standardized neurologic examination (all administered by specially trained nurses), and a 1-hour field battery of neuropsychological tests. All subjects who were capable completed a 10-minute cognitive screening instrument, the 3MS examination (54). The majority (84%) of subjects were subsequently examined by a board-certified gerontopsychiatrist and underwent neuroimaging. All data were then reviewed at a consensus diagnostic conference that included senior neuropsychologists and cognitive neuroscientists, gerontopsychiatrists, and board-certified neurologists. The year of disease onset was assigned retrospectively as the point at which subjects unambiguously met DSM-III-R criteria for dementia.

Subjects and Classification

Dementia was diagnosed in the 5092 study subjects by means of DSM-III-R criteria. AD diagnoses were based on NINCDS-ADRDA criteria (56), and VaD diagnoses were established by following NINDS-AIREN criteria (57, 58). Other diagnoses were made using standardized research criteria (50). A mild/ambiguous category was applied to those subjects who evidenced MCI or borderline impairment, suggestive of incipient AD, but who did not meet threshold criteria for specific dementia (52).

Ultimately, 335 cases of dementia were identified, of which 180 patients had MR imaging studies suitable for quantitative MR analysis. The various diagnostic categories to which the subjects were assigned are listed in Table 1. To simplify the parametric analyses, these categories were abridged to four classification groups: AD, MCI, clinical disorder not AD (including all disorders other than AD and MCI), and normal (including the 24 control subjects who underwent *APOE* genotyping, MR imaging, and the full assessment process). Subjects with AD constituted the largest group, followed by MCI and VaD subjects (see Table 1). Since $\epsilon 4$ has been associated with all three of these diagnostic categories (59–62), for several of the analyses, AD, VaD, and MCI subjects were combined into a single group and analyzed by presence or absence of the $\epsilon 4$ allele.

Because the influence of *APOE* may be apparent early in the evolution of AD, a finding substantiated in the Cache County population (50), one might expect to observe a differential distribution of *APOE* genotypes in demented subjects of varying ages. Furthermore, because the initial analysis of this population showed *APOE* effects related to AD and VaD subjects (22, 50, 62), AD and VaD subjects were combined in some analyses to increase the sample size and statistical power. Similarly, the MCI subjects were added to this analysis, because some had prodromal AD (63).

MR Imaging

MR imaging was performed at a regional medical center using a 0.5-T scanner with a quadrature head coil. The following imaging procedures were used: sagittal scans were T1-weighted with parameters of 500/15/2 (TR/TE/excitations), an acquisition matrix of 256×256 , a field of view of 24 cm, and a section thickness of 5 mm with a 1-mm gap. Axial intermediate (proton density-weighted) and T2-weighted spin-echo images were acquired with parameters of 3148/31/90/1, a field of view of 22 cm, a matrix of 256×256 , and a section

TABLE 1: Study demographics

	No.	Men	Women	Mean Age (SD)	Apolipoprotein E Genotype	
					ε 4 ⁻	ε 4 ⁺
Control group	24	11	13	77 (7.6)	15	9
Alzheimer disease	90	33	57	83 (6.8)	25	65
Mild cognitive impairment	39	20	19	83 (6.6)	12	27
Vascular dementia	21	8	13	83 (7.8)	15	6
Cerebrovascular disease	2	0	2	80 (2.1)	0	2
Frontal lobe dementia	3	1	2	81 (3.2)	1	2
Parkinson's disease	4	3	1	77 (3.9)	3	1
Psychiatric disorder	2	2	0	77 (15.6)	2	0
Dementia unknown	15	7	8	81 (7.4)	8	7
Alcohol dementia	2	2	0	84 (9.9)	2	0
Amnesic syndrome	2	0	2	80 (9.2)	1	1

thickness of 5 mm with a 1.5-mm gap. Coronal images obtained with a dual spin-echo technique (3046/30;90/1) were 3-mm thick with a 0.3-mm gap, a 22-cm field of view, and a matrix of 256×240 .

Quantitative MR Imaging

Quantitative analyses were performed using well-established image analysis protocols, which have been published previously (11, 12, 43). Briefly, quantification was accomplished through multispectral segmentation of the proton density- and T2-weighted images, in which white and gray matter and CSF were separated from one other using ANALYZE software (64). Application of such multispectral segmentation permitted the volumetric quantification of the following structures: total intracranial volume (TICV); total brain volume; total brain CSF; and total ventricular, temporal horn, lateral ventricle, white matter, gray matter, and hippocampal volumes. The VBR was calculated on the basis of total ventricular and brain volume. With the exception of hippocampal volume, which was obtained in the coronal plane, all quantitative analyses were based on the axial images. All quantitative MR analyses were performed blinded to subjects' age, sex, *APOE* genotype, and diagnostic classification. The TICV measure was used to control for head size variation, except in VBR (11). Because the only MR scanner in Cache County was a 0.5-T unit and all our previous research had relied on images acquired at 1.5T on a different machine, we scanned four patients on both machines with comparable imaging protocols. We then subjected the digital data to identical multispectral image analyses, comparing the segmented images from the two systems. Raters had initially established ANALYZE intraclass reliability ($r > 0.9$) on images from the 1.5-T magnet, except for white and gray matter volumes, for which reliability was less ($r > 0.7$). These same raters found similar reliability coefficients when analyzing the subjects imaged on the two machines. Segmentation routines for white versus gray matter differentiation have some inherent variation based on operator classification. Since the contrast between brain parenchyma, CSF, and bone provides clear MR signal demarcation, the quantitative MR measures dependent on these tissue/compartments segmentations were more reliable (ie, TICV, total brain volume, CSF, and ventricular volumes). However, for whole-brain white and gray matter volumes, slight differences of even a pixel width result in variability when attempting to differentiate these tissues with this routine (65). Thus, although excellent inter- and intrarater reliabilities that exceeded $r = 0.9$ were achieved for TICV, VBR, whole brain, hippocampal volume, and all CSF and ventricular measures, the coefficients were approximately $r = 0.7$ for estimates of white and gray matter volume.

Statistical Analysis

Owing to the multiple diagnostic categories and quantitative MR measures, the analysis of *APOE* genotype was simplified by combining ε4 homozygotes (ε4/4) and heterozygotes (ε4/ε3) into a single group, designated as ε4+, because a single copy of ε4 was sufficient to produce the association of ε4 to AD in this population (50). This type of *APOE* classification into ε4+ versus ε4- has been used by others (36, 37, 63). The nine subjects who were ε2/ε4 were classified as ε4+, even though some evidence suggests that ε2 may be protective (66, 67). However, no such protection influence was seen in the Cache County cohort (50) nor in this study when ε2/ε4 subjects were included or excluded. Thus, quantitative MR findings were analyzed by *APOE* status (ε4+ versus ε4-) and four levels of diagnostic classification (control group, AD, MCI, and clinical disorder not AD; see Table 2). As mentioned earlier, in some of the analyses, AD, VaD, and MCI subjects were pooled together in a single group. The justification for including the MCI subjects was based on the fact that they do display mild cognitive deficits and this condition may be a prodrome for dementia (52, 63, 68). Furthermore, combining all subjects with various levels of cognitive impairment provides a single group with a continuum of cognitive deficits so that quantitative MR findings may be compared by presence or absence of ε4. In every case in which these three groups were combined, they were first analyzed separately. Separate analysis yielded no significant findings. Head size was controlled by using TICV as a covariate in analyses of covariance, which allowed for heterogeneous regression slopes among groups (ANCOVET[†] 69). Other covariates included age at assessment and time since onset of dementia (when present) as a measure of LOD. Since the hippocampal volume calculations relied on separate images obtained in the coronal plane, there were two fewer hippocampal volumetric analyses than available for other structures, which were all obtained in the axial plane. Temporal horn and hippocampal volumes were calculated separately for left and right as well as a combined mean for each. In all, 17 quantitative analyses were performed. For simplicity in data analysis and presentation, we chose the following three representative measures for graphic depiction of test results: total brain volume (brain), total hippocampal volume (hippocampus), and VBR.

Results

Demographics and Descriptive Findings: Quantitative MR × Diagnosis × *APOE* Classification

Table 1 summarizes group composition by sex and *APOE* classification. Graphic depiction of the

TABLE 2: Quantitative MR findings in brain parenchyma

Sam- ple Size	Control Group		Alzheimer's Disease		Mild Cognitive Impairment		Clinical Disorder, Not Alzheimer's Disease	
	€ 4 ⁻ n = 15, Mean (SD)	€ 4 ⁺ n = 9, Mean (SD)	€ 4 ⁻ n = 25, Mean (SD)	€ 4 ⁺ n = 65, Mean (SD)	€ 4 ⁻ n = 12, Mean (SD)	€ 4 ⁺ n = 27, Mean (SD)	€ 4 ⁻ n = 32, Mean (SD)	€ 4 ⁺ n = 19, Mean (SD)
Age*	76.73 (8.06)	78.89 (6.55)	85.32 (6.88)	81.86 (6.46)	82.92 (6.57)	72.19 (6.82)	82.06 (8.00)	79.63 (6.14)
GM	676.08 (68.50)	710.15 (59.20)	592.91 (84.03)	568.60 (89.64)	581.26 (127.20)	579.03 (76.43)	553.27 (111.54)	549.61 (109.94)
WM	473.99 (45.83)	457.45 (58.47)	434.33 (105.20)	393.90 (107.65)	429.49 (71.02)	442.92 (128.28)	454.32 (130.81)	457.61 (115.18)
Brain	1150.07 (102.51)	1167.61 (101.70)	1027.23 (136.55)	962.50 (120.04)	1010.75 (137.09)	1021.95 (108.82)	1007.58 (135.66)	1007.22 (161.75)
TICV	1380.26 (128.90)	1460.78 (137.86)	1392.35 (146.53)	1316.79 (124.70)	1353.84 (148.29)	1334.98 (120.66)	1359.03 (167.59)	1348.36 (175.85)
VBR	3.70 (2.08)	4.25 (2.20)	7.11 (3.07)

Note.—GM indicates gray matter; WM, white matter; brain, total brain volume; TICV, total intracranial volume; VBR, ventricle-to-brain ratio.

* Age at time of clinical assessment.

mean (\pm SD) for whole brain and hippocampal volumes along with VBR is presented in Figure 1 for each diagnostic category. As expected, control subjects had the highest brain volume, with the exception of one outlier with Parkinson's disease and two subjects who suffered from alcoholism. Likewise, VBR was smallest in control subjects, except for the person with Parkinson's disease. In contrast, hippocampal volume did not exhibit such a clear separation across groups. Two subjects with frontal lobe dementia and the subject with Parkinson disease had the smallest hippocampal volume, followed by the AD subjects. In regard to *APOE* effects, generally, €4+ subjects had smaller hippocampal volumes, even among the control group. The smallest brain volumes were observed in AD and VaD subjects with at least one €4 allele. However, none of the €4-related effects remained significant after controlling for age.

Table 2 provides the quantitative MR results across the four broad diagnostic groups for each morphometric measure (Table 3). Comparing groups by diagnostic classification yielded the expected results, wherein the control subjects had significantly less atrophy than the others ($F_{\text{brain}} = 6.13$, $df = 3, 184$, $P = .001$; $F_{\text{VBR}} = 3.26$, $df = 3, 184$, $P = .012$; $F_{\text{hippocampus}} = 3.59$, $df = 3, 178$, $P = .008$). As with Figure 1, inspection of Table 2 reveals that, in many comparisons, subjects with an €4 allele had signs of greater atrophy; however, no significant *APOE* effects, either primary or interactive, were observed when age was controlled. On most quantitative MR measures, subjects in the AD, MCI, and clinical disorder not AD groups had significantly more atrophic changes than did control subjects, but they did not differ from one another.

Age, Quantitative MR Findings, and *APOE*

For the analyses that follow, AD, VaD, and MCI subjects were combined, and Figure 2 shows scatter plots representing each data point by subjects' age for brain and hippocampal volumes and VBR. Each panel of the figure also shows two regression lines, representing €4+ and €4- subjects, respectively. Inspection of Figure 2B shows that for €4+ subjects, hippocampal volume was smallest in those between the ages of 65 and 75 years. The difference was significant ($t = 3.92$, $df = 39$, $P = .0001$). In addition, €4+ subjects had slightly smaller total brain volume by age, although this difference was not significant. VBR was also greater in €4+ subjects at younger ages, but two distinct outliers with severe cerebral atrophy were largely responsible for this finding (see Fig 2C). Identical analyses were performed by separating AD, VaD, and MCI subjects into separate groups. When examined independently, no significant findings were observed.

TABLE 3: Brain CSF quantitative MR findings

Sample Size	Control Group		Alzheimer's Disease		Mild Cognitive Impairment		Clinical Disorder, Not Alzheimer's Disease	
	ϵ_{4-} n = 15 Mean (SD)	ϵ_{4+} n = 9 Mean (SD)	ϵ_{4-} n = 25 Mean (SD)	ϵ_{4+} n = 65 Mean (SD)	ϵ_{4-} n = 12 Mean (SD)	ϵ_{4+} n = 27 Mean (SD)	ϵ_{4-} n = 32 Mean (SD)	ϵ_{4+} n = 19 Mean (SD)
CSF	230.19 (86.90)	293.16 (72.66)	365.12 (76.27)	354.30 (82.37)	343.09 (64.39)	313.03 (74.61)	351.45 (99.93)	341.14 (108.85)
IV	1.67 (0.63)	1.58 (0.32)	1.76 (0.62)	1.56 (0.80)	1.40 (0.48)	1.57 (0.75)	1.49 (0.71)	1.66 (0.51)
RTH	0.30 (0.36)	0.61 (0.80)	1.38 (1.37)	1.67 (1.74)	1.36 (1.42)	0.75 (0.79)	1.16 (0.82)	1.22 (0.99)
LTH	0.47 (0.45)	0.62 (0.92)	1.35 (1.20)	1.51 (1.44)	1.26 (1.64)	0.69 (0.62)	0.99 (1.05)	1.49 (1.25)
HORN	0.39 (0.37)	0.62 (0.84)	1.37 (1.21)	1.59 (1.54)	1.31 (1.52)	0.72 (0.66)	1.08 (0.86)	1.35 (1.08)
III	2.12 (0.88)	2.01 (0.74)	2.66 (0.83)	2.40 (0.82)	2.57 (1.05)	2.07 (0.89)	2.47 (0.95)	2.39 (0.81)
LAT	37.76 (21.72)	44.11 (20.89)	65.20 (29.72)	67.03 (32.61)	58.17 (30.01)	55.48 (31.47)	68.74 (32.92)	63.56 (38.13)
TVENT	42.32 (22.65)	48.94 (21.85)	72.35 (32.04)	74.17 (35.47)	64.76 (33.75)	60.55 (33.46)	74.85 (34.79)	70.31 (40.24)
SUBCSF	187.87 (69.31)	244.23 (56.80)	292.76 (59.53)	280.13 (67.20)	278.34 (58.81)	252.48 (57.09)	276.60 (79.19)	270.83 (92.10)

Note.—CSF indicates whole brain cerebrospinal fluid; IV, fourth ventricle; RTH, right temporal horn; LTH, left temporal horn; III, third ventricle; LAT, lateral ventricle; BRAIN, total brain; TVENT, total ventricular volume; SUBCSF, subarachnoid CSF; HORN, average temporal horn volume; HIPP, average hippocampal volume.

LOD, Quantitative MR Findings, APOE, and Diagnostic Classification

LOD was estimated for AD and VaD subjects. Combining AD and VaD subjects, Figure 3 offers a graphic summation of LOD effects in relation to total brain and hippocampal volumes and VBR by ϵ_{4} status. For those subjects with disease duration of less than 1 year, hippocampal volume was significantly smaller in ϵ_{4+} subjects ($t = 2.84$, $df = 6$, $P = .015$). All other comparisons and analyses yielded no significant *APOE* by LOD interaction effects. When AD and VaD subjects were assessed separately, no significant *APOE* \times LOD effect was observed.

Cognitive Performance, Quantitative MR Findings, and APOE

To examine the relationship between 3MS examination performance and quantitative MR findings across the four major diagnostic groups, 3MS scores were divided into three groups: ≥ 91 (normal range), 71 to 90 (mild to moderate impairment), and ≤ 70 (moderate to severe impairment) to provide three levels of cognitive impairment for comparison. Results are summarized in Figure 4. In the majority of comparisons, 3MS performance was inversely related to degree of atrophy. Significant 3MS main effects on total brain volume ($F = 14.27$, $df = 2$, 188, $P = .0001$) and hippocampal volume ($F = 4.04$, $df = 2$, 182, $P = .001$) along with VBR ($F = 12.67$, $df = 2$, 188, $P = .001$) were found. However, there were no significant *APOE* effects.

Discussion

Our population-based study of dementia provides a quantitative description of brain morphology among subjects 65 years and older, all with known *APOE* genotype. As expected, quantitative MR findings showed distinct morphometric differences between healthy control subjects and those with dementia. Dementia, regardless of origin, was generally associated with demonstrable atrophy manifested by smaller brain and hippocampal volumes, along with larger ventricular volume. These observations are consistent with other studies (35, 36, 38–42). Level of cognitive impairment, as measured by 3MS performance, was inversely related to degree of atrophy. Of particular interest in this context was the group with ambiguous impairment or MCI. MCI subjects may be prodromal for AD (63), but at the time of initial assessment they did not meet criteria for diagnosis of AD (or any other dementing illness). As a group, the MCI subjects displayed global atrophic indicators intermediate to those observed in AD or VaD subjects, but distinctly more than those in control subjects. Unlike the AD and VaD subjects, however, the MCI group had hippocampal volumes similar to that of control subjects. For MCI subjects, it may be that structural





FIG 1. A–C, Graphic depictions of mean (\pm SD) for brain volume (A), hippocampal volume (B), and VBR (C) for $\epsilon 4+$ and $\epsilon 4-$ subjects for each diagnostic classification: control group, Alzheimer disease (AD), mild ambiguous (M/A), vascular dementia (VaD), cerebrovascular disease (CVA), frontal lobe dementia (FLA), Parkinson's disease (Parkinson), psychiatric disorder (Psych), dementia unknown (Dem Unk), alcoholism (ETOH), and amnesic disorder (Mem Amn). The first numeric value represents the total number of $\epsilon 4-$ subjects, with the second numeric value indicating sample size of the $\epsilon 4+$ subjects.



integrity of the hippocampus, even in the context of more global brain atrophy (elevated VBR and decreased brain volume), delays the frank expression of dementia. Recently, Jack et al (63) found that MCI patients who premorbidly had smaller hippocampal volume were more likely to progress to an AD classification (35, 70). Undoubtedly, such subjects are imbedded in our MCI population, and, on follow-up, will most likely meet criteria for AD (such an investigation in this population is currently underway). Taken together, these findings suggest that atrophy is a nonspecific effect of essentially all age-related disorders that may have a component of cognitive impairment or of actual dementia (71, 72). The intactness of the hippocampus may be critical in preserving cognitive status during the preclinical phase of AD.

As anticipated, age effects were apparent in this population of elderly subjects, who ranged from 65 to more than 100 years old (14, 35, 49, 73–75). Increased age was associated with decreased brain and hippocampal volumes along with increased

VBR. For hippocampal volume, $\epsilon 4+$ subjects had smaller hippocampi than did $\epsilon 4-$ subjects if they were under the age of 75. Over that age, no significant $\epsilon 4$ effect was observed. Likewise, $\epsilon 4+$ subjects who were early in the disease process (ie, 1 year or less) had the smallest hippocampal volumes. Since the AD risk effect of *APOE* may be mediated by age (76, 77), the morphometric change associated with presence of an $\epsilon 4$ allele may be expressed at a younger age and possibly in the mildest stage of illness. Bondi et al (78), Caselli et al (79), and Smith et al (80) found possession of an $\epsilon 4$ allele to be associated with cognitive impairment in nondemented subjects. Plassman et al (43), Reiman et al (8), and Tohgi et al (44) have all found smaller hippocampal volumes in cognitively normal $\epsilon 4+$ individuals. These observations would fit with an early rather than late expression of an *APOE* effect on brain morphology. Clearly, even if there is an early $\epsilon 4$ effect on brain morphology, the current study suggests that once the AD or VaD progresses to full expression or the AD




FIG 2. A–C, Scatter plots for subjects with AD and combined MCI and VaD by age for brain volume (A), hippocampal volume (B), and VBR (C) based on the following correlations: Brain volume, $r_{\epsilon 4-} = -.26$, $P \leq .01$; $r_{\epsilon 4+} = -.19$, $P \leq .01$; hippocampal volume, $r_{\epsilon 4-} = 0.17$, $P \leq .05$; $r_{\epsilon 4+} = 0.01$, $P > .05$; and VBR, $r_{\epsilon 4-} = -.34$, $P \leq .01$; $r_{\epsilon 4+} = 0.13$, $P \leq .05$.

or VaD subject is beyond 75 years of age, *APOE* genotype probably does not systematically influence gross brain morphology. Potentially, this is an important finding of this investigation, as it confirms other studies that did not detect an $\epsilon 4$ effect on brain morphology with either advanced age or disease severity, yet raises the possibility of detecting morphologic differences earlier in life or in the disease process. Obviously, additional research is needed.

The standard limitations of the quantitative MR technique always are a concern in data interpretation (65) and must be taken into consideration when reviewing these observations. For example, the lack of effect may be due to measurement error or to insensitivity of the quantitative MR technique in discriminating hippocampal disease at the microstructural level (65). For instance, the segmentation method used to assess hippocampal volume is merely a global measure of hippocampal anatomy. While volume is related to neuronal cell count

(81), it is nonetheless a crude measure of hippocampal integrity. Thus, the real proof of any *APOE* effect at the hippocampal level (or within any other brain structure) must await histologic examination or additional sensitive neuroimaging analyses of the hippocampus; eg, with high-resolution MR imaging, functional MR imaging, or spectroscopy (82, 83).

Another limitation is that the current study was not longitudinal. Since presence of $\epsilon 4$ confers increased risk and earlier onset of AD, any potential *APOE* effect on brain morphology may only be seen in the earliest stage of the disease (maybe even prodromally) and may only be discovered by tracking quantitative MR findings in the at-risk population from prodromal stages through diagnosis of dementia. Such longitudinal investigations combined with regional analyses (4, 5, 9, 83–86), in particular, hippocampal volume in the context of other temporal lobe morphologic measures (63, 72, 83), may be critical in establishing any *APOE* ef-





FIG 3. A–C, Mean (\pm SD) for brain volume (A), hippocampal volume (B), and VBR (C) by length of disease (LOD) in years.




fects that can be determined by neuroimaging analysis.


Like other epidemiologic studies of dementia, the Cache County study shows an enhanced risk and younger age of AD onset in subjects who possess at least one copy of the $\epsilon 4$ allele (50). Several lines of evidence point to a role of *APOE* in neuronal maintenance and repair (77, 87, 88), and a number of studies have shown that the presence of the $\epsilon 4$ allele is associated with development of β -amyloid and neurofibrillary tangles (24, 89–91). Additionally, the $\epsilon 4$ allele at *APOE* may be associated with vascular angiopathy (30) and ischemic vascular disease (92); hence, the relationship between *APOE* $\epsilon 4$ and VaD. Heuristically, the presence of the $\epsilon 4$ allele, if it is associated with some form of impaired neuronal maintenance, could be a pathologic factor associated with the development of AD and VaD (59). However, while this as well as other epidemiologic studies have shown age-related *APOE* effects, more than half the Cache

County population of elderly who had an $\epsilon 4$ allele did not develop AD by age 100. Obviously, other factors in addition to *APOE* genotype participate in the development of AD (27, 28, 31), such as other susceptibility genes (93) or environmental modifiers (eg, head injury) (94, 95). The overriding finding of this investigation is that once dementia is expressed, *APOE* genotype does not appear to play a unique or major role in the degree of cerebral atrophy.

Yasuda et al (20) made an interesting observation in a cross-sectional, clinical sample of AD subjects recruited from a hospital dementia service in Japan. While all subjects with AD had significantly reduced brain volume as compared with control subjects, AD subjects with $\epsilon 4/4$ genotype actually had the least degree of brain atrophy. These investigators interpreted this observation to mean that $\epsilon 4+$ subjects incurred the disease early and, owing to their relatively young age, the ravages of a degenerative disease interacting with age had not




A



B

FIG 4. A–C, Mean (\pm SD) for brain volume (A), hippocampal volume (B), and VBR (C) for 3MS performance, which has been segregated into three levels: normal (> 91), mild to moderate impairment (71–90), and moderate to severe impairment (< 70). Correlation values (Spearman rank) for each brain measure by 3MS performance level are shown in the boxes.



C

reached their peak. Hence, there was less atrophy in these younger $\epsilon 4+$ AD subjects. Yasuda et al used a sample of convenience, and the mean age of their $\epsilon 4/4$ subjects was 70 years. In contrast, the $\epsilon 4+$ group in the Cache County study, while younger (see Table 1) than the $\epsilon 4-$ subjects, were considerably older (mean age, 80+ years) than the subjects studied by Yasuda et al (20). Although symptom duration was reported, the volumetric findings in the study by Yasuda et al were not analyzed by disease duration. Jack et al (5), controlling for disease duration, found no difference in hippocampal volume between $\epsilon 4+$ and $\epsilon 4-$ AD subjects.

Conclusion

We found generalized atrophy to be ubiquitous across all forms of dementia and associated disorders investigated in this population-based study. A subtle *APOE* $\epsilon 4$ effect may be present early in the AD and/or VaD disease process, wherein quantitative MR analysis of hippocampal volume shows

greater hippocampal atrophy. However, once an individual is advanced in age or in progression of disease, any *APOE* effect on gross brain morphology is mitigated. Accordingly, there appear to be no lasting morphologic effects detected by quantitative MR imaging that are associated with presence or absence of the $\epsilon 4$ allele. This observation is consistent with other recent reports (36, 37). In addition, degree of brain atrophy is inversely related to 3MS performance, irrespective of *APOE* genotype.

Acknowledgments

Grateful acknowledgement is given to Drs. John C. S. Breitner, Bonita Wyse, Kathleen A. Welsh-Bohmer, JoAnn Tschanz, and Brenda Plassman. The technical assistance and expertise of Tracy Abildskov is greatly appreciated, as is the secretarial assistance of Johanna Carlquist and Allicen Neal.

References

1. Cummings JL, Vinters HV, Cole GM, Khachaturian ZS. **Alzheimer's disease: etiologies, pathophysiology, cognitive reserve, and treatment opportunities.** *Neurology* 1998;51:S2–S17

2. Jorm AF, Jolley D. **The incidence of dementia: a meta-analysis.** *Neurology* 1998;51:728-733
3. Mayeux R, Saunders AM, Shea S, et al. **Utility of the apolipoprotein E genotype in the diagnosis of Alzheimer's disease.** *N Engl J Med* 1998;338:506-511
4. Kidron D, Black SE, Stanchev P, et al. **Quantitative MR volumetry in Alzheimer's disease.** *Neurology* 1997;49:1504-1512
5. Jack CR Jr, Petersen RC, Xu YC, et al. **Hippocampal atrophy and apolipoprotein E genotype are independently associated with Alzheimer's disease.** *Ann Neurol* 1998;43:303-310
6. Ylikoski R, Ylikoski A, Erkinjuntti T, et al. **Differences in neuropsychological functioning associated with age, education, neurological status, and magnetic resonance imaging findings in neurologically healthy elderly individuals.** *Appl Neuropsychol* 1998;5:1-14
7. Insausti R, Juottonen K, Soininen H, et al. **MR volumetric analysis of the human entorhinal, perirhinal, and temporopolar cortices.** *AJNR Am J Neuroradiol* 1998;19:659-671
8. Reiman EM, Uecker A, Caselli RJ, et al. **Hippocampal volumes in cognitively normal persons at genetic risk for Alzheimer's disease.** *Ann Neurol* 1998;44:288-291
9. Kohler S, Black SE, Sinden M, et al. **Memory impairments associated with hippocampal versus parahippocampal-gyrus atrophy: an MR volumetry study in Alzheimer's disease.** *Neuropsychologia* 1998;36:901-914
10. Kaye JA. **Diagnostic challenges in dementia.** *Neurology* 1998;51:S45-S52
11. Blatter DD, Bigler ED, Gale SD, et al. **Quantitative volumetric analysis of brain MR: normative database spanning five decades of life.** *AJNR Am J Neuroradiol* 1995;16:241-251
12. Bigler ED, Blatter DD, Anderson CV, et al. **Hippocampal volume in normal aging and traumatic brain injury.** *AJNR Am J Neuroradiol* 1997;18:11-23
13. Skoog I, Palmertz B, Andreasson LA. **The prevalence of white-matter lesions on computed tomography of the brain in demented and nondemented 85-year-olds.** *Psychiatry Neurol* 1994;7:169-175
14. Mueller EA, Moore MM, Kerr DCR, et al. **Brain volume preserved in healthy elderly through the eleventh decade.** *Neurology* 1998;51:1555-1562
15. Libon DJ, Bogdanoff B, Cloud BS, et al. **Declarative and procedural learning, quantitative measures of the hippocampus, and subcortical white alterations in Alzheimer's disease and ischaemic vascular dementia.** *J Clin Exp Neuropsychol* 1998;20:30-41
16. Skoog I, Berg S, Johansson B, Palmertz B, Andreasson LA. **The influence of white matter lesions on neuropsychological functioning in demented and non-demented 85-year-olds.** *Acta Neurol Scand* 1996;93:142-148
17. Skoog I, Hesse C, Fredman P, Andreasson L, Palmertz B, Blennow K. **Apolipoprotein E in cerebrospinal fluid in 85-year-old subjects.** *Arch Neurol* 1997;54:267-272
18. Price DL, Sisodia SS, Borchelt DR. **Alzheimer disease: when and why?** *Nat Genet* 1998;19:314-316
19. Martinez M, Campion D, Brice A, et al. **Apolipoprotein E $\epsilon 4$ allele and familial aggregation of Alzheimer disease.** *Arch Neurol* 1998;55:810-816
20. Yasuda M, Mori E, Kitagaki H, et al. **Apolipoprotein E $\epsilon 4$ allele and whole brain atrophy in late-onset Alzheimer's disease.** *Am J Psychiatry* 1998;155:779-784
21. Tilvis RS, Strandberg TE, Juva K. **Apolipoprotein E phenotypes, dementia, and mortality in a prospective population sample.** *J Am Geriatr Soc* 1998;46:712-715
22. Kuller LH, Shemanski L, Manolio T, et al. **Relationship between ApoE, MRI findings, and cognitive function in the cardiovascular health study.** *Stroke* 1998;29:388-398
23. Breitner JCS, Jarvik GP, Plassman BL, Saunders AM, Welsh KA. **Risk of Alzheimer disease with the $\epsilon 4$ allele for apolipoprotein E in a population-based study of men aged 62-73 years.** *Alzheimer Dis Assoc Disord* 1998;12:40-44
24. Warzok RW, Kessler C, Apel G, et al. **Apolipoprotein $\epsilon 4$ promotes incipient Alzheimer pathology in the elderly.** *Alzheimer Dis Assoc Disord* 1998;12:33-39
25. Berg L, McKeel DW, Miller JP, et al. **Clinicopathologic studies in cognitively healthy aging and Alzheimer disease.** *Arch Neurol* 1998;55:326-335
26. Jonker C, Schmand B, Lindeboom J, Havekes LM, Launer LJ. **Association between apolipoprotein E $\epsilon 4$ and the rate of cognitive decline in community-dwelling elderly individuals with and without dementia.** *Arch Neurol* 1998;55:1065-1069
27. Blacker D, Wilcox MA, Laird NM, et al. **Alpha-2 macroglobulin is genetically associated with Alzheimer disease.** *Nat Genet* 1998;19:357-360
28. Wu WS, Holmans P, Wavrant-DeVrieze F, et al. **Genetic studies on chromosome 12 in late-onset Alzheimer disease.** *JAMA* 1998;280:619-622
29. Meyer MR, Tschanz JT, Norton MC, et al. **APOE genotype predicts when-not whether-one is predisposed to develop Alzheimer disease.** *Nat Genet* 1998;19:321-322
30. Katzman R, Zhang MY, Chen PJ, et al. **Effects of apolipoprotein E on dementia and aging in the Shanghai survey of dementia.** *Neurology* 1997;49:779-785
31. Rogaeva E, Premkumar S, Song Y, et al. **Evidence for an Alzheimer disease susceptibility locus on chromosome 12 and for further locus heterogeneity.** *JAMA* 1998;280:614-618
32. Saunders AM, Hulette C, Welsh-Bohmer KA, et al. **Specificity, sensitivity, and predictive value of apolipoprotein-E genotyping for sporadic Alzheimer's disease.** *Lancet* 1996;19:224-228
33. Strittmatter WJ, Roses AD. **Apolipoprotein E and Alzheimer's disease.** *Annu Rev Neurosci* 1996;19:53-77
34. Bigler ED, Lowry CM, Anderson CV, et al. **A population based study of APOE, Alzheimer's disease and quantitative magnetic resonance imaging.** *J Int Neuropsychol Soc* 1997;3:241
35. Jack CR, Petersen RC, Xu Y, et al. **Rate of medial temporal lobe atrophy in typical aging and Alzheimer's disease.** *Neurology* 1998;51:993-999
36. Barber R, Gholkar A, Scheltens P, et al. **Apolipoprotein E $\epsilon 4$ allele, temporal lobe atrophy, and white matter lesions in late-life dementias.** *Arch Neurol* 1999;56:961-965
37. Barber R, Ballard C, McKeith IG, Gholkar A, O'Brien JT. **MRI volumetric study of dementia with Lewy bodies: a comparison with AD and vascular dementia.** *Am Acad Neurol* 2000;54:1304-1309
38. Lehtovirta M, Soinene H, Laasko MP, et al. **SPECT and MRI analysis in Alzheimer's disease: relation to apolipoprotein E $\epsilon 4$ allele.** *J Neurol Neurosurg Psychiatry* 1996;60:644-649
39. Lehtovirta M, Laasko MP, Soininen H, et al. **Volumes of hippocampus, amygdala and frontal lobe in Alzheimer patients with different apolipoprotein E genotypes.** *Neuroscience* 1995;67:65-72
40. Soininen H, Partanen K, Pitkanen A, et al. **Decreased hippocampal volume asymmetry on MRIs in nondemented elderly subjects carrying the apolipoprotein E $\epsilon 4$ allele.** *Neurology* 1995;45:391-392
41. Soininen HS, Riekkinen PJ. **Apolipoprotein E, memory and Alzheimer's disease.** *Trends Neurosci* 1996;19:224-228
42. Geroldi C, Pihlajamaki M, Laasko MP, et al. **APOE- $\epsilon 4$ is associated with less frontal and more medial temporal lobe atrophy in AD.** *Neurology* 1999;53:1825-1832
43. Plassman BL, Welsh-Bohmer KA, Bigler ED, et al. **Apolipoprotein E $\epsilon 4$ and hippocampal volume in twins with normal cognition.** *Neurology* 1997;48:985-989
44. Tohogi H, Takahashi S, Kato E, et al. **Reduced size of right hippocampus in 39- to 80-year-old normal subjects carrying the apolipoprotein E $\epsilon 4$ allele.** *Neurosci Lett* 1998;236:21-24
45. Hirano N, Mori E, Yasuda M, et al. **Lack of association of apolipoprotein E $\epsilon 4$ allele dose with cerebral glucose metabolism in Alzheimer disease.** *Alzheimer Dis Assoc Disord* 1998;12:362-367
46. Van Dyck CH, Gelernter J, MacAvoy MG. **Absence of an apolipoprotein E $\epsilon 4$ allele is associated with increased parietal regional cerebral blood flow asymmetry in Alzheimer disease.** *Arch Neurol* 1998;1460-1466
47. Reiman EM, Caselli RJ, Yun LS. **Preclinical evidence of Alzheimer's disease in persons homozygous for the $\epsilon 4$ allele for apolipoprotein E.** *N Engl J Med* 1996;334:752-758
48. Small GW, Mazziotta JC, Collins MT. **Apolipoprotein E type 4 allele and cerebral glucose metabolism in relatives at risk for familial Alzheimer's disease.** *JAMA* 1995;273:942-947
49. Smith CD, Malcein M, Meurer K, Schmitt FA, Markesbery WR, Pettigrew LC. **MRI temporal lobe volume measures and neuropsychological function in Alzheimer's disease.** *J Neuroimaging* 1999;9:2-9
50. Breitner JCS, Wyse BW, Anthony JC, et al. **APOE- $\epsilon 4$ count predicts age when prevalence of AD increases, then declines: the Cache County study.** *Neurology* 1999;53:321-331
51. Tschanz JT, Welsh-Bohmer KA, Skoog I, et al. **Dementia diagnoses from clinical and neuropsychological data compared: the Cache County study.** *Am Acad Neurol* 2000;54:1290-1296

52. Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. **Mild cognitive impairment: clinical characterization and outcome.** *Arch Neurol* 1999;30:303–308
53. Saunders AM, Strittmatter WJ, Schmechel D, et al. **Association of apolipoprotein E allele $\epsilon 4$ with late onset familial and sporadic Alzheimer's disease.** *Neurology* 1993;43:1467–1472
54. Teng EL, Chui HC. **The modified Mini-Mental State (3MS) Examination.** *J Can Psychiatry* 1987;48:314–318
55. Khachaturian AS, Gallo JJ, Breitner JCS. **Performance characteristics of a two-stage dementia screen in a population sample.** *J Clin Epidemiol* (in press)
56. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. **Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA work group under the auspices of department of Health and Human Services task force on Alzheimer's disease.** *Neurology* 1984;939–944
57. Roman GC, Tatemichi TK, Erkinjuntti T, et al. **Vascular dementia: diagnostic criteria for research studies: report of the NINDS-AIREN International Workshop.** *Neurology* 1993;43:243–245
58. Tatemichi TK, Sacktor N, Mayeux R. **Dementia Associated with Cerebrovascular Disease, Other Degenerative Diseases, and Metabolic Disorders.** New York: Raven; 1994
59. Haan MN, Shemanski L, Jagust WJ, Manolio TA, Kuller L. **The role of APOE $\epsilon 4$ in modulating effects of other risk factors for cognitive decline in elderly persons.** *JAMA* 1999;282:40–46
60. Tierney MC, Szalai JP, Snow WG, et al. **A prospective study of the clinical utility of ApoE genotype in the prediction of outcome in patients with memory impairment.** *Am Acad Neurol* 1996;45:149–153
61. Dik MG, Jonker C, Bouter LM, Geerlings MI, Van Kamp GJ, Deeg DJH. **APOE- $\epsilon 4$ is associated with memory decline in cognitively impaired elderly.** *Neurology* 2000;
62. Carmelli D, Swan GE, Reed T, et al. **Midlife cardiovascular risk factors, ApoE, and cognitive decline in elderly male twins.** *Neurology* 1998;50:1580–1585
63. Jack CR, Petersen RC, Xu YC, et al. **Prediction of AD with MRI-based hippocampal volume in mild cognitive impairment.** *Neurology* 1999;52:1397–1403
64. Robb R. **Three-dimensional Biomedical Imaging.** New York: VCH Publishers; 1995
65. Jack CR, Theodore WH, Cook M, McCarthy G. **MRI-based hippocampal volumetrics: data acquisition, normal ranges, and optimal protocol.** *Magn Reson Imaging* 1995;13:1057–1064
66. Corder EH, Saunders AM, Risch NJ, et al. **Protective effect of apolipoprotein E type 2 allele for late onset Alzheimer's disease.** *Nat Genet* 1994;7:180–184
67. Lai F, Kamman E, Rebeck GW, Anderson A, Chen Y, Nixon RA. **APOE genotype and gender effects on Alzheimer disease in 100 adults with Down syndrome.** *Neurology* 1999;53:331–336
68. Friedrich MJ. **Mild cognitive impairment raises Alzheimer disease risk.** *JAMA* 1999;282:621–622
69. Maxwell SE, Delaney HD. **Designing Experiments and Analyzing Data: A Model Comparison Perspective.** Belmont, CA: Wadsworth; 1990
70. Juottonen K, Laakso MP, Partanen K, Soininen H. **Comparative MR analysis of the entorhinal cortex and hippocampus in diagnosing Alzheimer disease.** *AJNR Am J Neuroradiol* 1999;20:139–144
71. Zakzanis KK. **Quantitative evidence for neuroanatomic and neuropsychological markers in dementia of the Alzheimer's type.** *J Clin Exp Neuropsychol* 1998;20:259–269
72. Frisoni GB, Laakso MP, Beltramello A, et al. **Hippocampal and entorhinal cortex atrophy in frontotemporal dementia and Alzheimer's disease.** *Neurology* 1999;52:91–100
73. Raz N. **Aging of the brain and its impact on cognitive performance: integration of structural and functional findings.** In: Craik FIM, Salthouse TA, eds. *Aging of the Brain and Its Impact on Cognitive Performance: Integration of Structural and Functional Findings.* Mahwah, NJ: Erlbaum; 2000 (in press)
74. Bertoni MA, Sclavi NE, Sauer HJ. **Volumetry of the hippocampus and amygdala with magnetic imaging.** *Int J Neuroradiol* 1998;4:291–295
75. Mu Q, Xie J, Wen Z, Weng Y, Shuyun Z. **A quantitative MR study of the hippocampal formation, the amygdala, and the temporal horn of the lateral ventricle in healthy subjects 40 to 90 years of age.** *AJNR Am J Neuroradiol* 1999;20:207–211
76. Blacker D, Haines JL, Rodes L, et al. **ApoE-4 and age at onset of Alzheimer's disease: the NIMH genetics initiative.** *Neurology* 1997;48:139–147
77. Masliah E, Mallory M, Veinbergs I, Miller A, Samuel W. **Alterations in apolipoprotein E expression during aging and neurodegeneration.** *Prog Neurobiol* 1996;50:493–503
78. Bondi MW, Salmon DP, Monsch AU. **Episodic memory changes are associated with the apoE- $\epsilon 4$ allele in nondemented older adults.** *Neurology* 1995;45:2203–2206
79. Caselli RJ, Graff-Radford NR, Reiman EM, et al. **Preclinical memory decline in cognitively normal apolipoprotein E- $\epsilon 4$ homozygotes.** *Neurology* 1999;53:201–207
80. Smith GE, Bohac DL, Waring SC, et al. **Apolipoprotein E genotype influences cognitive "phenotype" in patients with Alzheimer's disease but not in healthy control subjects.** *Neurology* 1998;50:355–362
81. Harding AJ, Halliday GM, Kriil JJ. **Variation in hippocampal neuron number with age and brain volume.** *Cereb Cortex* 1998;8:710–718
82. Friedman SD, Brooks WM, Jung RE, Hart BL, Yeo RA. **Proton MR spectroscopic findings correspond to neuropsychological function in traumatic brain injury.** *AJNR Am J Neuroradiol* 1998;19:1879–1885
83. Small SA, Petera GM, DeLaPaz R, Mayeux R, Stern Y. **Differential dysfunction of the hippocampal formation among elderly with memory decline and Alzheimer's disease.** *Ann Neurol* 1999;45:466–472
84. Thompson PM, Moussai J, Zohoori S, et al. **Cortical variability and asymmetry in normal aging and Alzheimer's disease.** *Cereb Cortex* 1998;8:492–509
85. Pfefferbaum A, Sullivan EV, Jernigan TL, et al. **A quantitative analysis of CT and cognitive measures in normal aging and Alzheimer's disease.** *Psychiatry Res* 1990;35:115–136
86. Pfefferbaum A, Sullivan E, Rosenbloom MJ, Mathalon DH, Lim KO. **A controlled study of cortical gray matter and ventricular changes in alcoholic men over a 5-year interval.** *Arch Gen Psychiatry* 1998;55:905–912
87. Poirier J, Hess M, May PC, Finch CE. **Astrocytic apolipoprotein E mRNA and GFAP mRNA in hippocampus after entorhinal cortex lesioning.** *Mol Brain Res* 1991;11:97–106
88. Poirier J. **Apolipoprotein E in animal models of CNS injury and in Alzheimer's disease.** *Trends Neurosci* 1994;17:525–530
89. Schmechel DE, Saunders AM, Strittmatter WJ, et al. **Increased amyloid B-peptide deposition in cerebral cortex as a consequence of apolipoprotein E genotype in late-onset Alzheimer's disease.** *Neurobiology* 1993;90:9749–9753
90. Gomez-Isla T, West HL, Rebeck W, et al. **Clinical and pathological correlates of apolipoprotein E $\epsilon 4$ in Alzheimer's disease.** *Ann Neurol* 1996;39:62–70
91. Polvikoski T, Sulkava R, Haltia M, et al. **Apolipoprotein E, dementia, and cortical deposition of B-amyloid protein.** *N Engl J Med* 1996;333:1242–1247
92. McCarron MO, Delong D, Alberts MJ. **APOE genotype as a risk factor for ischemic cerebrovascular disease.** *Neurology* 1999;53:1308–1311
93. Pericak-Vance MA, Bebout JL, Gaskell PJ, et al. **Linkage studies in familial Alzheimer's disease: evidence for chromosome 19 linkage.** *Am J Hum Genet* 1991;48:1034–1050
94. Mayeux R, Ottman R, Maestre G, et al. **Synergistic effects of traumatic head injury and apolipoprotein- $\epsilon 4$ in patients with Alzheimer's disease.** *Neurology* 1995;45:555–557
95. Breitner JCS, Gau BA, Welsh KA, et al. **Inverse association of anti-inflammatory treatments and Alzheimer's disease: initial results of a co-twin control study.** *Neurology* 1994;44:227–232