



Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Primary adenocarcinoma of the middle ear.

W Adam, J C Johnson, D J Paul, K Clausen and D E Schuller

AJNR Am J Neuroradiol 1982, 3 (6) 674-676

<http://www.ajnr.org/content/3/6/674.citation>

This information is current as
of August 10, 2025.

Primary Adenocarcinoma of the Middle Ear

Wynn Adam,¹ John C. Johnson,¹ David J. Paul,¹ Kathryn Clausen,² and David E. Schuller³

Adenocarcinoma of the middle ear is a rare condition. It should be considered in the differential diagnosis of lytic lesions of the temporal bone particularly in a patient with a history of chronic ear infection. The following case illustrates the possible extensive radiographic changes produced by this disease.

Case Report

A 38-year-old woman was referred for evaluation and treatment of recurrent drainage and pain in the right ear. She also had a several month history of mild dysphagia and change in her voice.

Her problems began 22 years before when she developed recurrent episodes of right otitis media, commonly associated with otorrhea. A progressive ipsilateral facial paralysis developed. Attempted decompression of the right facial nerve 18 years before admission resulted in its sacrifice. A facial suspension was performed but the symptoms progressed. A mastoidectomy resulted in the pathologic diagnosis of a benign congenital cyst of the petrous part of the temporal bone. Four years before admission granulation tissue was removed through her ear. The pathologic diagnosis was again a simple cyst. Four months before admission, bloody otorrhea and dysphagia developed. Because of the severity and persistence of her symptoms, she was admitted to the hospital for further evaluation.

The skull, mastoid, and polytome examinations revealed a destructive lesion of the petrous part of the right temporal bone with a sharply margined peripheral bony rim (fig. 1A). Tumor calcification was not seen within the lesion which appeared to expand the bone. A contrast-enhanced computed tomographic (CT) scan showed a ringlike enhancing lesion in an expanded destroyed part of the right temporal bone (fig. 1B). A nonenhanced scan was not obtained. This lesion extended into the middle cranial fossa and the anterior part of the posterior fossa with displacement of the third, fourth, and lateral ventricles to the left (fig. 1C). Angiography disclosed a highly vascular tumor supplied primarily by meningeal branches of the carotid and vertebral-basilar circulations (figs. 2A and 2B).

The patient underwent surgical resection of a large vascular neoplasm with a blood loss of 20 U during surgery. The excised tumor was bulky and hemorrhagic. There were several reddish brown irregular masses weighing 36 g altogether. Microscopically the tumor was unusual. It was composed of highly vascularized, colloid-filled follicles lined by a single layer of plump, clear cells. In many areas a papillary growth pattern was evident (fig. 2C). This

pattern resulted in a close histologic resemblance to thyroid carcinoma. Tumor calcification was not present. The final pathologic diagnosis was adenocarcinoma of the middle ear.

Comparison of this lesion with the tissue excised 18 years previously showed a few points of similarity. That specimen contained cystic spaces lined by simple cuboidal cells and a few of the cells had clear cytoplasm. This tended to confirm that the tumor was primary in the temporal bone and not metastatic from some other site. This also supported the slowly evolving nature of this neoplasm.

A follow-up CT scan 5 months after surgery showed a postsurgical defect with no evidence of recurrent tumor. The ventricular system had returned to a normal configuration.

Discussion

Adenocarcinoma of the middle ear is a rare lesion generally not diagnosed early in its course. Fewer than 25 cases have been reported [1–8]. Typically, the patients have a long history of otitis media with otorrhea, otalgia, and hearing loss. Other associated but less common problems are facial paralysis, vertigo, tinnitus, fullness in the ear, headache, and retroauricular swelling [1, 3, 4].

The radiographic manifestations are nonspecific and include clouding of the mastoid antrum and air cells, bone destruction, and a soft-tissue mass in the tympanic cavity [3–5]. CT changes apparently have not been described.

The differential diagnosis for a destructive lesion of the temporal bone is extensive [9, 10]. Primary neoplasms include squamous cell carcinoma [11], adenoid cystic carcinoma, adenocarcinoma, osteosarcoma, chondrosarcoma, rhabdomyosarcoma [12], paraganglionoma [13], hemangioma, facial nerve neuroma, meningioma [13, 14], teratoma, giant cell tumor, and hemangiopericytoma [15]. Secondary malignancies may produce bony changes by local invasion, by blood-borne metastases [16] (particularly from breast, lung, and kidney), and by involvement with hematologic malignancies such as multiple myeloma or leukemia. Other destructive processes of the temporal bone are cholesteatoma, histiocytosis X [17], malignant otitis externa, tuberculosis, Wegener granulomatosis, surgical defects, mucocele [18], and cholesterol granuloma [19]. The slow

Received September 15, 1981; accepted after revision February 16, 1982.

¹Department of Radiology, Ohio State University Hospital, 410 W. 10th Ave., Columbus, OH 43210. Address reprint requests to W. Adam.

²Department of Pathology, Ohio State University Hospital, Columbus, OH 43210.

³Department of Otolaryngology, Ohio State University Hospital, Columbus, OH 43210.

AJNR 3:674–676, November/December 1982 0195–6108/82/0306–0674 \$00.00 © American Roentgen Ray Society

Fig. 1.—**A**, Towne view of skull. Large lytic lesion (*short arrows*) in right petrous bone. Tumor displaces normal bone as it advances along its superior margin (*long arrow*). **B**, Contrast-enhanced axial CT scan. Both destruction and expansion of right petrous bone. Posterior cortex of petrous bone displaced posteriorly and medially (*arrow*). **C**, Axial CT image 1 cm higher than **B**. Growth of tumor into temporal lobe and posteriorly into cerebellum with displacement of fourth ventricle (*arrow*) to left.





Fig. 2.—**A**, Lateral subtraction view from selective right external carotid arteriogram. Tumor supplied by multiple branches of external carotid artery and displays intense vascular stain (*arrow*). **B**, Lateral subtraction view of selective right vertebral arteriogram. Inferior part of tumor supplied by men-

ingeal branches of vertebral artery. Both intense tumor blush (*long arrows*) and early draining vein (*short arrow*) apparent. **C**, Colloid-filled follicles and papillary growth pattern of middle ear tumor (H and E $\times 100$).

growth characteristics and vascularity of this destructive lesion exclude most of these considerations and limit the differential diagnosis to: paraganglioma, hemangioma, hemangiopericytoma, slowly growing vascular metastasis, giant cell tumor, and primary adenocarcinoma of the middle ear [4].

The histopathology of this case is interesting. A previous report of two cases of adenocarcinoma of the middle ear described trabecular and gland-forming patterns [1]. We found only one previous report of papillary adenocarcinoma of the middle ear [2]. This combination of papillary and follicular pattern may be confused with metastatic tumor. The preoperative CT and angiographic findings in our case suggested a destructive, vascular neoplasm. Preoperative

transcatheter embolization of the feeding arteries might have been beneficial. Stone et al. [4] described a similar large intraoperative blood loss.

Primary adenocarcinoma of the middle ear should be included in the diagnostic considerations of a destructive lesion of the temporal bone, particularly if the clinical course has been lengthy.

REFERENCES

1. Fayemi AO, Toker C. Primary adenocarcinoma of the middle ear. *Arch Otolaryngol* 1975;101:449-452
2. Siedentop HK, Jeantet C. Primary adenocarcinoma of the middle ear: report of three cases. *Ann Otol Rhinol Laryngol*

- 1961;70:719-733
3. Dehner LP, Chen KTK. Primary tumors of the external and middle ear: benign and malignant glandular neoplasms. *Arch Otolaryngol* 1980;106:13-19
 4. Stone HE, Lipa M, Bell RD. Primary adenocarcinoma of the middle ear. *Arch Otolaryngol* 1975;101:702-705
 5. Derlacki EL, Barney PL. Adenomatous tumors of the middle ear and mastoid. *Laryngoscope* 1976;86:1123-1135
 6. Wagenfeld DJH, Keane T, Van Nostrand AWP, Bryce DP. Primary carcinoma involving the temporal bone: analysis of twenty-five cases. *Laryngoscope* 1980;90:912-919
 7. Sinha PP, Aziz HI. Treatment of carcinoma of the middle ear. *Radiology* 1978;126:485-487
 8. Conley J, Schuller DE. Malignancies of the ear. *Laryngoscope* 1976;86:1147-1163
 9. Livingston PA. Differential diagnosis of radiolucent lesions of the temporal bone. *Radiol Clin North Am* 1974;12:571-583
 10. Scully RE, Galdalsini JJ, McNeely BU. Case records of the Massachusetts General Hospital—weekly clinicopathological exercises. *N Engl J Med* 1980;302:456-460
 11. Wang CC. Radiation therapy in the management of carcinoma of the external auditory canal, middle ear, or mastoid. *Radiology* 1975;116:713-715
 12. Dehner LP, Chen KTK. Primary tumors of the external and middle ear: a clinicopathologic study of embryonal rhabdomyosarcoma. *Arch Otolaryngol* 1978;104:399-403
 13. Chen KTK, Dehner LP. Primary tumors of the external and middle ear: a clinicopathologic study of 14 paragangliomas and 3 meningiomas. *Arch Otolaryngol* 1978;104:253-359
 14. Guzowski J, Paparella MM, Rao KN, Hoshino T. Meningiomas of the temporal bone. *Laryngoscope* 1976;86:1141-1146
 15. Osborne DR, Dubois P, Drayer B, Sage M, Burger P, Heinz ER. Primary intracranial meningeal and spinal hemangiopericytoma: radiologic manifestations. *AJNR* 1981;2:69-74
 16. Stucker FJ, Holmes WF. Metastatic disease of the temporal bone. *Laryngoscope* 1976;86:1136-1140
 17. McCaffrey TV, McDonald TJ. Histiocytosis X of the ear and temporal bone: review of 22 cases. *Laryngoscope* 1979;89:1735-1742
 18. Osborn AG, Parkin JL. Mucocoele of the petrous temporal bone. *AJR* 1979;132:680-681
 19. Nager GT, Vanderveen TS. Cholesterol granuloma involving the temporal bone. *Ann Otol* 1976;85:204-209