



Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents

**FRESENIUS
KABI**

[WATCH VIDEO](#)

AJNR

Fractional Flow on TOF-MRA as a Measure of Stroke Risk in Children with Intracranial Arterial Stenosis

A.Y. Ibrahim, A. Amirabadi, M.M. Shroff, N. Dlamini, P. Dirks and P. Muthusami

This information is current as of August 16, 2025.

AJNR Am J Neuroradiol 2020, 41 (3) 535-541

doi: <https://doi.org/10.3174/ajnr.A6441>

<http://www.ajnr.org/content/41/3/535>

Fractional Flow on TOF-MRA as a Measure of Stroke Risk in Children with Intracranial Arterial Stenosis

 A.Y. Ibrahim,  A. Amirabadi,  M.M. Shroff,  N. Dlamini,  P. Dirks, and  P. Muthusami

ABSTRACT

BACKGROUND AND PURPOSE: Conventional angiography is the criterion standard for measuring intracranial arterial stenosis. We evaluated signal intensity ratios from TOF-MRA as a measure of intracranial stenosis and infarct risk in pediatric stroke.

MATERIALS AND METHODS: A retrospective study was undertaken in children with intracranial arterial stenosis, who had TOF-MRA and conventional angiography performed within 6 months. Arterial diameters were measured for percentage stenosis. ROI analysis on TOF-MRA measured signal intensity in pre- and poststenotic segments, with post-/pre-signal intensity ratios calculated. The Pearson correlation was used to compare percentage stenosis on MRA with conventional angiography and signal intensity ratios with percentage stenosis; the point-biserial correlation was used for infarcts compared with percentage stenosis and signal intensity ratios. Sensitivity, specificity, and positive and negative predictive values were calculated for determining severe ($\geq 70\%$) stenosis from MRA and signal intensity ratios against the criterion standard conventional angiography. $P < .05$ was considered statistically significant.

RESULTS: Seventy stenotic segments were found in 48 studies in 41 children (median age, 11.0 years; range, 5 months to 17.0 years; male/female ratio, 22:19): 20/41 (48.8%) bilateral, 11/41 (26.8%) right, and 10/41 (24.4%) left, with the most common site being the proximal middle cerebral artery (22/70, 31%). Moyamoya disease accounted for 27/41 (65.9%). Signal intensity ratios and conventional angiography stenosis showed a moderate negative correlation ($R = -0.54$, $P < .001$). Receiver operating characteristic statistics showed an area under the curve of 0.86 for using post-/pre-signal intensity ratios to determine severe ($\geq 70\%$) carotid stenosis, yielding a threshold of 1.00. Sensitivity, specificity, and positive and negative predictive values for severe stenosis were the following—MRA: 42.8%, 58.8%, 30.0%, and 71.4%; signal intensity ratio > 1.00 : 97.1%, 77.8%, 71.7%, and 97.4%; combination: 75.5%, 100%, 100%, and 76.8%, respectively. Signal intensity ratios decreased with increasing grade of stenosis (none/mild-moderate/severe/complete, $P < .001$) and were less when associated with infarcts (0.81 ± 0.52 for arteries associated with downstream infarcts versus 1.31 ± 0.55 for arteries without associated infarcts, $P < .001$).

CONCLUSIONS: Signal intensity ratios from TOF-MRA can serve as a noninvasive measure of intracranial arterial stenosis and allow identification of high-risk lesions in pediatric stroke.

ABBREVIATIONS: CA = conventional angiography; SI = signal intensity

Stroke in children is 1 of the top 10 causes of death in childhood.^{1,2} The incidence of all strokes in children is reported to be 2–13 per 100,000 person-years in developed countries.^{3–6}

Received August 16, 2019; accepted after revision January 4, 2020.

From the Department of Diagnostic Imaging (A.Y.I., A.A., M.M.S., P.M.), Division of Neurology (N.D.) and Division of Neurosurgery (P.D.), Department of Surgery, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada; and Department of Clinical Sciences (A.Y.I.), Faculty of Medicine, Yarmouk University, Irbid, Jordan.

The study was approved by the Institutional Research Ethics Board with REB 1000062860. Written informed consent was obtained from all patients/parents prior to the procedures.

Please address correspondence to Ala' Y. Ibrahim, MD, Department of Clinical Sciences, Faculty of Medicine, Yarmouk University, P.O. Box 566, 21163 Irbid, Jordan; e-mail: dr.alaa_yonis@yahoo.com; @Ala_Y_Ibrahim

<http://dx.doi.org/10.3174/ajnr.A6441>

Childhood arteriopathies are the most common identifiable cause of arterial ischemic stroke, accounting for up to two-thirds of cases in a previously healthy child.^{7–11} The most common causes of intracranial arterial steno-occlusive disease in children are Moyamoya disease, focal dissection, transient cerebral arteriopathy, and post-varicella arteriopathy,^{12–14} though a number of other syndromic and nonsyndromic associations exist.¹⁵

Stroke in children is commonly assessed and followed by MR imaging, which is noninvasive and free of ionizing radiation.¹² Vascular assessment is commonly performed using TOF-MRA, which gives a global assessment of vascular disease. However, TOF-MRA is amenable to several artifacts resulting from the inherent technique and is known to overestimate stenotic lesions while underevaluating arteries that are severely affected.^{16–18}




FIG 1. Measurement of arterial signal intensity from TOF-MRA using ROI analysis. A, Prestenotic ROI. B, Poststenotic ROI.

Often, for further clarification of equivocal findings, contrast-enhanced MR imaging or CT angiography is performed, though these too have certain drawbacks. Conventional angiography (CA) remains the criterion standard test to diagnose intracranial arterial stenosis and accurately quantify severity.^{19–21} This is, however, an invasive test, using ionizing radiation and requires general anesthesia in most children and a dedicated pediatric neuroangiographic service to maximize safety.

The signal intensity (SI) within arteries on TOF-MRA is dependent on the number of moving spins and the velocity of blood flow and thus has the potential to noninvasively assess the downstream effect of arterial stenosis. Ratios derived from SI on TOF-MRA have been shown to correlate with cerebral perfusion as measured by CT perfusion and SPECT in adults with intracranial carotid stenosis.^{22,23} It is not known whether arterial SI ratios measured from TOF-MRA can accurately estimate the degree of intracranial carotid stenosis as measured on conventional angiography in children presenting with stroke and whether these ratios can help identify high-risk lesions. We hypothesized the following: 1) SI ratios calculated from TOF-MRA are significantly correlated with percentage stenosis measured from conventional angiography, 2) SI ratios calculated from TOF-MRA are significantly correlated with infarct burden, and 3) SI ratios serve as stronger predictors of severe (>70%) intracranial arterial stenosis than stenosis measured from TOF-MRA.

MATERIALS AND METHODS

Subjects and Imaging

An institutional review board–approved retrospective study was undertaken on children 18 years of age or younger with an imaging diagnosis of intracranial carotid artery stenosis at our tertiary pediatric institution (The Hospital for Sick Children) between January 1, 2000, and November 10, 2018. The need for individual consent for this retrospective review was waived by the Institutional Research Ethics Board. A keyword search was performed on the radiology departmental data base to identify all studies reported with intracranial arterial stenosis. Studies were included that had both TOF-MRA and conventional angiography performed within 6 months of each other. Patients with tandem lesions (ie, 2 noncontiguous stenotic lesions), those with MRA and conventional angiography performed more than 6 months apart, and patients with suboptimal imaging quality resulting in unreliable SI or stenosis measurement were excluded.

MR imaging was performed on a 1.5T magnet (Achieva; Philips Healthcare, Best, the Netherlands) or a 3T magnet (Magnetom Skyra; Siemens, Erlangen, Germany, or Achieva) with a 32-channel transmit-receive head coil. The examination included standard T1-, T2-, and FLAIR-weighted sequences, followed by TOF-MRA with the following parameters—1.5T MRA: TR/TE = 25/6.91 ms, flip angle = 20°, bandwidth = 109 Hz, FOV = 18 cm, spatial resolution = 0.70 × 0.90 × 1.0 mm

(interpolated to 0.5 × 0.5 × 0.5 mm), 1-mm section thickness with 0.5-mm overlap; 3T MRA: TR/TE = 25/3.81 ms, flip angle = 18°, bandwidth = 186 Hz, FOV = 18 cm, spatial resolution = 0.40 × 0.70 × 1.0 mm (interpolated to 0.5 × 0.5 × 0.5 mm), 1.0-mm section thickness with 0.5-mm overlap. Conventional angiography was performed by 1 of 2 interventional neuroradiologists (P.M. or M.M.S.) in a biplane neuroangiography suite (Artis Q; Siemens). All procedures were performed with the patient under general anesthesia through femoral artery access and systemic heparinization using a 4F diagnostic catheter with standard projections and an iodinated contrast agent (iohexol, Omnipaque 300 mg I/mL; GE Healthcare, Piscataway, New Jersey) injected through a power injector at 3–5 mL/s, depending on the age and arterial size.

Data Analysis and Image Processing

Images and reports were analyzed by a fellowship-trained pediatric neuroradiologist (A.Y.I.), with discrepancies resolved by the senior author (P.M.). Electronic chart review and image analysis on the PACS system were performed for included patients to collect the following data: demographics (age at diagnosis, sex), time interval between MRA and conventional angiography, laterality of stenosis (right/left/bilateral), specific location/segment of artery involved (petrous ICA, cavernous ICA, paraclinoid ICA, pre-communicating ICA, terminal ICA, postbifurcation), any ischemic events regardless of the association with the territory of the stenotic lesion, and any ischemic or hemorrhagic events not associated with the stenotic lesion. Stenosis was regarded as symptomatic if ischemic lesions on MR imaging were territorially consistent with the stenotic lesion. Downstream infarcts were defined as any abnormal bright T2/FLAIR lesions in the vascular distribution of the stenotic segment without mass effect.

Arterial diameters were measured to assess the percentage of ICA stenosis on MRA and CA. The percentage stenosis on MRA and catheter cerebral angiography was measured as described previously.²⁴ In summary, percentage stenosis of an intracranial artery is $[(1 - (D_{\text{stenosis}}/D_{\text{normal}}))] \times 100$, where D_{stenosis} is the diameter of the ICA at the maximally stenotic segment and D_{normal} is the diameter of the most adjacent proximal normal artery (usually the petrous ICA). If the proximal segment was diseased, contingency sites were chosen to measure D_{normal} : the most adjacent nondiseased distal artery (second choice, usually the terminal ICA); if the entire intracranial artery was diseased, the most distal, parallel, nontortuous normal segment was measured (third choice). ROI analysis was performed to measure intravascular SI




FIG 2. Flow chart showing patient selection for our study. SIR indicates signal intensity ratios.

Table 1: Sites of intracranial arterial stenosis in our cohort (n = 70 stenotic segments)

Sites	
Right	14
Left	12
Bilateral	22
Associated with infarct	36/70 (51.4%)
Specific location	
Cervical and petrous	4 (6%)
Cavernous	3 (4%)
Postclinoid	20 (29%)
Postcommunicating and terminal	21 (30%)
Postterminal (A1 ACA or M1 MCA)	22 (31%)

Note:—ACA indicates anterior cerebral artery.

Table 2: Underlying diagnosis in our cohort of children with stroke and intracranial arterial stenosis (n = 41)

Clinical Diagnosis	No.
Moyamoya disease	15
Neurofibromatosis type 1 with Moyamoya	6
Trisomy-21 with Moyamoya	3
Dissection	2
Primary CNS vasculitis	3
Varicella vasculitis	2
TCA	3
Sickle cell disease with Moyamoya	3
Other	
Systemic large vessel vasculitis	1
Hemolytic uremic syndrome	1
Thalassemia with Moyamoya	1
Hurler syndrome	1

Note:—TCA indicates transient cerebral arteriopathy.

from TOF-MRA. ROIs were manually drawn on TOF-MRA axial source images (Fig 1). The upper cervical ICA SI was measured at approximately the C2 vertebral level for the prestenotic segment (henceforth, SI-ICA-pre) and from the MCA for the poststenotic segment (henceforth, SI-ICA-post). Each SI-ICA value was measured as the mean of 2 ROIs drawn on adjacent axial sections or adjacent arterial segments with no overlap of ROIs. Ratios of SI were calculated, as SI-post/SI-pre-stenosis.

Statistical Analysis

Categorical variables were expressed as number and percentage; continuous variables were expressed as median and range. For statistical purposes, stenosis was graded as follows: none (0%), mild (<50%), moderate (50%–69%), severe (70%–99%), and complete (100%), as previously described.²⁴ Correlations of percentage stenosis on MRA with percentage stenosis on conventional angiography and with SI ratios were performed using the Pearson correlation; the presence of infarcts and the development of new infarcts were compared with percentage stenosis and SI ratios using a point-biserial correlation. Receiver operating characteristic curve analysis was performed on SI ratios as a determinant

of percentage stenosis (from CA), from which the area under the curve and threshold value were obtained. Sensitivity, specificity, positive predictive value, and negative predictive value for determining severe ($\geq 70\%$) stenosis by MRA and SI ratios were calculated, with conventional angiography percentage stenosis as the criterion standard. A P value < .05 was considered statistically significant. All analyses were performed with SPSS (Version 25.0; IBM, Armonk, New York).

RESULTS

Of 55 children with intracranial ICA stenosis, 41 met the inclusion criteria, which resulted in 48 studies with 70 stenotic arteries and 36 associated downstream arterial infarcts. Figure 2 shows the flow chart for patient selection. The median age of our cohort was 11 years (range, 5 months to 17 years). Of these, 19/41 patients (46.3%) were girls. MR imaging was performed at 1.5T in 25/48 and 3T in 23/48 studies. Bilateral stenosis was seen in 20/41 patients (48.8%); 11/41 patients (26.8%) had stenosis on the right side only and 10/41 patients (24.4%) had stenosis on the left side only. The most common site for stenosis (Table 1) was distal to the ICA bifurcation (22/70, 31%), (ie, the MCA), followed by the terminal ICA (21/70, 30%). The most common underlying diagnosis was Moyamoya disease (alone or in association with neurofibromatosis 1, Trisomy-21, sickle cell disease, or thalassemia) accounting for nearly two-thirds of patients (27/41, 65.9%). Other diagnoses are shown in Table 2. Associated downstream infarcts were seen in 36/48 studies (75%), 24/48 (50%) had nonspecific white matter lesions, and 4/48 (8.3%) had hemorrhage.

The mean interval between MR imaging and CA was 22.7 ± 32.7 days (range, 0–123 days). Severe stenosis ($\geq 70\%$) was detected by MRA in 50/70 segments (71.4%) and by conventional angiography in 46/70 segments (65.7%) (Table 3). Although there was excellent correlation between MRA and conventional angiographic stenosis (Pearson $R = 0.93$, $P < .001$), MRA had a sensitivity, specificity, positive predictive value, and negative predictive value of 42.8%, 58.8%, 30.0%, and 71.4%, respectively,

for severe ($>70\%$) stenosis. The severity of stenosis on MRA correlated with the infarct burden ($r_{pb} = 0.44$, $P < .001$). Figure 3 shows 2 relevant sample cases.

Mean ratios of SI (post/pre) were as follows: 1.58 ± 0.51 (no stenosis), 1.40 ± 0.50 (mild-moderate stenosis, $<70\%$), 0.90 ± 0.48 (severe stenosis, $70\%–99\%$), and 0.60 ± 0.33 (complete, 100%), with a significant difference of means by 1-way ANOVA ($P < .001$). A significant moderate negative correlation was seen between post-/prestenosis SI ratios and percentage stenosis on conventional angiography ($R = -0.54$, $P < .001$). Receiver operating characteristic statistics (Fig 4) showed an area under the curve of 0.86 (95% CI, 0.78–0.94) for using post-/pre-ICA SI ratios to determine severe ($\geq 70\%$) stenosis. An SI ratio of 1.00 was determined to be the threshold for obtaining maximum sensitivity and specificity for severe stenosis. By means of this SI ratio cutoff value alone and in combination with MRA percentage stenosis, the sensitivity, specificity, positive predictive value, and negative predictive value for determining severe ($>70\%$) stenosis were 97.1%, 77.8%, 71.7%, 97.4%, and 75.5%, 100%, 100%, and 76.8%, respectively.

Table 3: ICA stenosis severity, comparison of MRA with catheter angiography evaluations ($n = 96$ arteries: 70 stenotic and 26 nonstenotic)

% Stenosis	Arteries on MRA (No.)	Arteries on CA (No.)
≤ 50	31	29
51–69	16	21
70–99	4	32
100	46	14




FIG 3. A–C, A 15-month-old boy with intracranial vasculopathy and bilateral intracranial arterial stenosis. A, Maximum-intensity-projection reformat from TOF-MRA shows bilateral terminal ICA stenosis, measured on MRA as right: 100%; left: 86%. SI ratios were measured as right: 1.0; left: 1.3. B, The child presented with extensive MCA territory infarct. C, MR imaging performed 5 years later. Axial FLAIR image shows no new infarcts in the left cerebral hemisphere. D–F, A 15-year-old boy with right-sided intracranial arteriopathy. D, Maximum-intensity-projection reformat from TOF-MRA shows 60% stenosis of the supraclinoid right ICA, with additional mild narrowing of the proximal A1 anterior cerebral artery and a normal appearance of the left ICA. The SI ratios were measured as right: 0.60; left: 1.3. E, The patient presented with acute infarction of the right basal ganglia. F, MRA performed 22 months later shows near-complete resolution of the supraclinoid ICA stenosis, consistent with transient cerebral arteriopathy.

There was a moderate negative correlation between the presence of infarcts and SI ratios ($r_{pb} = -0.4$, $P < .001$). The mean post-/pre-SI-ratio for arteries associated with downstream infarcts was 0.81 ± 0.52 , whereas for arteries without associated infarcts, it was 1.31 ± 0.55 , the difference being significant by an unpaired t test ($P < .001$).

DISCUSSION

There is a strong relationship between impaired hemodynamics related to intracranial arterial stenosis and increased stroke risk.¹⁹ The severity of intracranial arterial stenosis and its progression with time dictate treatment protocols and clinical decision-making, including the decision to surgically revascularize. Conventional angiography is the criterion standard for the diagnosis of the presence and severity of intracranial arterial stenosis.¹⁹ Noninvasive MR angiography and CTA have partially replaced conventional angiography, which has up to a 1% risk of stroke, a 4% risk of TIA, and nearly a 1% mortality rate.^{25–27} Additionally, in children, there are the added concerns with ionizing radiation and general anesthesia. Most institutions worldwide have now accepted MRA as the optimal technique for following pediatric patients with stroke over the course of their disease. However, measuring the severity of mild-moderate stenosis and detecting severe stenosis ($>70\%$) by TOF-MRA in children is fraught with challenges resulting from artifacts inherent to the TOF technique. These factors, as well as technical differences among scans make interpretation of the progression of arterial stenosis across time an imprecise science. Basing disease severity on SI alone would also imply that the physiologic state at the time of scanning would be an important consideration. More

recent versions of TOF-MRA have focused on improving the spatial resolution and signal-to-noise ratios using overlapping slices (multiple overlapping thin-slab acquisition) or variable flip angles to maintain the relative spin saturations. How these affect SI in a diseased vessel is not fully known and can result in over- or underestimation of stenosis. To minimize the influence of acquisition parameters, we used the SI ratios instead. Signal intensity ratios on TOF-MRA have the potential to assess the fractional flow reserve of the assessed artery, thus serving as a noninvasive surrogate to identify high-risk intracranial stenosis, its severity, and functional significance.¹⁹

A recent evaluation of adult patients in the Stroke Outcomes and Neuroimaging of Intracranial Atherosclerosis (SONIA) and Warfarin-Aspirin Symptomatic Intracranial Disease (WASID) trials showed that decreased SI ratios across intracranial arterial stenoses correlated with misery perfusion to the downstream brain territories, thereby




FIG 4. Receiver operator characteristic curve for post-/prestenotic SI ratios to diagnose severe (>70%) intracranial arterial stenosis. The area under the curve was 0.86, with a determined cutoff of 1.00 (see text).

providing a simple and noninvasive predictor of high-risk intracranial stenosis.^{23,28} Signal intensities from TOF-MRA have also been shown to significantly correlate with end-diastolic flow velocities on duplex sonography performed 1 year following pial synangiosis in patients with symptomatic intracranial stenosis.²⁹ The reproducibility of SI ratios was shown in a recent study on atherosclerotic intracranial stenosis, with interobserver agreements of >80%.³⁰ In the present study, we add to this growing body of literature by showing that SI ratios derived from TOF-MRA in pediatric patients with stroke correlate with catheter angiographic severity of stenosis, serve as a strong discriminant for diagnosing severe (>70%) intracranial stenosis in these children, and are stronger predictors of infarct burden and risk than percentage stenosis.

Although there are studies comparing MRA and conventional angiography in various adult cohorts,^{31,32} this is, to the best of our knowledge, the first time this correlation is being reported in the pediatric stroke literature. The estimation of percentage stenosis for intracranial arteries from TOF-MRA is challenging, with the proximal arterial segments diminishing in size due to reduced flow, poststenotic dilation often being present, TOF-MRA artifacts in the stenotic and poststenotic segments, and a normal reduction in size after bifurcation. Selection of the segments used for this analysis is critical. As described above, we used a predefined method, choosing contingency sites if this was not possible. In our cohort of 41 children with 70 stenotic arterial segments, though we found a strong correlation ($R = 0.93$) between MRA and angiographic stenosis, we also showed that MRA was a poor discriminant of severe and mild-moderate intracranial stenosis. Serving as an estimate of the fractional flow

across the stenosis, SI ratios had higher sensitivity and specificity than percentage stenosis from MRA (97% and 78% versus 43% and 59%, respectively) and higher positive and negative predictive values (72% and 97% versus 30% and 71%, respectively). This finding is not dissimilar to the large body of literature regarding fractional flow reserve assessments in coronary artery disease, which illustrates that percentage stenosis is not an optimal marker for identifying disease of physiologic significance. Due to the invasiveness of intra-arterial wire measurements, noninvasive methods to measure fractional flow reserve have also been developed using coronary CT angiography.^{33,34}

While a similar severity of arterial stenosis may have different hemodynamic and functional significance in the cerebral circulation than it does in the coronary tree, it is conceivable that there are better predictors of stroke risk than the severity of stenosis as measured by percentage stenosis. Cerebrovascular reserve studies have shown that the measurement of cerebral blood flow is a strong predictor of stroke risk, independent of the severity of stenosis.³⁵ Several different methods have been used for this, including SPECT, PET, CT perfusion, MR imaging perfusion, and blood oxygen level-dependent MR imaging. A retrospective analysis of the WASID trial suggested that impaired collaterals on angiography may be a better predictor of stroke in the territory than percentage stenosis, suggesting a key role for impaired hemodynamics.^{22-24,28,36-38} This finding is not wholly unexpected, given that the intracranial circulation can effectively compensate for a stenotic lesion in a carotid artery by flow across the circle of Willis from the contralateral carotid artery and/or from the posterior circulation. Our finding of a stronger correlation of TOF-MRA-derived SI ratios than percentage stenosis with infarct burden implies that SI ratios could serve as a surrogate biomarker for impaired cerebrovascular reserve, independent of the severity of the stenosis. Liebeskind et al,²⁸ in a recent study, had similarly shown that SI ratios derived from TOF-MRA were a strong predictor of increased stroke risk in an adult cohort with intracranial atherosclerotic stenosis.

Our study had some limitations. The retrospective design of the study implied that all data were extracted from medical and imaging records. However, all images were analyzed and postprocessed by study authors using a standard method. Although our cohort included heterogeneous vascular pathologies, these are indicative of the diverse etiologies of pediatric stroke and retain their clinical relevance for neurologists and pediatricians involved in the care of children with stroke. Second, the use of MR imaging scanners of different strengths (1.5 and 3T) could result in some variability in TOF-MRA appearances and the measurement of stenosis. Again, however, the comparison with angiographic stenosis is useful information for clinical settings that commonly toggle between these scanners. Additionally, our use of SI ratios rather than SI itself is expected to negate scanner-related differences. Further research is required to analyze and compare SI ratio differences among scanners. Also, TOF-MRA has several artifacts that are known to affect stenosis measurements and could potentially affect SI values. We attempted to minimize this issue by measuring every value as the mean of 2 ROIs; Liebeskind et al,²⁸ using this method, had shown excellent inter-reader agreement.

Third, selecting an appropriate segment for placing an ROI for SI was challenging in several cases. Manual placement of the ROI, the level, and location chosen for measurement is subjective. We attempted to maintain consistency by acquiring measurements of arterial SI in the ICA at approximately the C2 vertebral level for the prestenotic segment and from at least 2 arterial segments downstream of the poststenotic segment. Last, intracranial flow measurements are subject to the immediate poststenotic flow effects being altered by the circle Of Willis, as well as by pial/dural collaterals.³⁹ The selection of the MCA as the site for measurement of poststenotic SI was based on this factor: This location would account for compensatory flow from the anterior and/or posterior communicating arteries into the poststenotic artery, for which direct measurements of percentage stenosis would not account.

CONCLUSIONS

SI ratios measured from TOF-MRA serve as a reliable estimate of the severity of intracranial ICA stenosis in pediatric stroke and can be used as a surrogate for noninvasively following the severity of stenosis across time. These ratios serve as fractional flow measurements, assessing the hemodynamic significance of stenosis and allowing identification of high-risk stenotic lesions. Future prospective studies in pediatric stroke, comparing these with more established cerebrovascular reserve methods, are required to further corroborate our findings.

ACKNOWLEDGMENTS

We would like to acknowledge the neuroradiologists and neurosurgeons at our institution and the Ontasian Imaging Lab office for their support through this project. We would also like to acknowledge the MR imaging, CT, and interventional radiology technologists who played a key role in the acquisition of images used in this study.

Disclosures: Manohar Shroff—UNRELATED: Expert testimony: Occasional medical expert on birth injury cases; Payment for lectures including service on speakers bureaus: Speakers bureau – one time for BioMarin, stipend for speaking at conference organized by Siemens Brazil – one time.

REFERENCES

- Heideman RL, Packer RJ, Albright LA, et al. Tumors of the central nervous system. In: Pizzo PA, ed. *Principles and Practice of Pediatric Oncology*. 3rd ed. Philadelphia: Lippencott; 1997:633–97
- National Center for Health Statistics. **Deaths, percentage of total deaths, and death rates for the 10 leading causes of death in selected age groups, by race and sex in the United States.** 2002. https://www.cdc.gov/nchs/data/nvsr/nvsr53/nvsr53_17.pdf. Accessed June 16, 2019
- Giroud M, Lemesle M, Madinier G, et al. **Stroke in children under 16 years of age: clinical and etiological difference with adults.** *Acta Neurol Scand* 1997;96:401–06 [CrossRef Medline](#)
- Lynch JK, Hirtz DG, DeVeber G, et al. **Report of the National Institute of Neurological Disorders and Stroke Workshop on Perinatal and Childhood Stroke.** *Pediatrics* 2002;109:116–23 [CrossRef Medline](#)
- Chung B, Wong V. **Pediatric stroke among Hong Kong Chinese subjects.** *Pediatrics* 2004;114:e206–12 [CrossRef Medline](#)
- Fullerton HJ, Wu YW, Zhao S, et al. **Risk of stroke in children: ethnic and gender disparities.** *Neurology* 2003;61:189–94 [CrossRef Medline](#)
- Fullerton HJ, Wu YW, Sidney S, et al. **Risk of recurrent childhood arterial ischemic stroke in a population-based cohort: the importance of cerebrovascular imaging.** *Pediatrics* 2007;119:495–501 [CrossRef Medline](#)
- Ganesan V, Prengler M, McShane MA, et al. **Investigation of risk factors in children with arterial ischemic stroke.** *Ann Neurol* 2003;53:167–73 [CrossRef Medline](#)
- Sträter R, Becker S, von Eckardstein A, et al. **Prospective assessment of risk factors for recurrent stroke during childhood: a 5-year follow-up study.** *Lancet* 2002;360:1540–05 [CrossRef Medline](#)
- Chabrier S, Husson B, Lasjaunias P, et al. **Stroke in childhood: outcome and recurrence risk by mechanism in 59 patients.** *J Child Neurol* 2000;15:290–94 [CrossRef Medline](#)
- Zimmer JA, Garg BP, Williams LS, et al. **Age-related variation in presenting signs of childhood arterial ischemic stroke.** *Pediatr Neurol* 2007;37:171–75 [CrossRef Medline](#)
- Wintermark M, Hills NK, DeVeber GA, et al; VIPS Investigators. **Clinical and imaging characteristics of arteriopathy subtypes in children with arterial ischemic stroke: results of the VIPS study.** *AJNR Am J Neuroradiol* 2017;38:2172–79 [CrossRef Medline](#)
- Sébire G, Fullerton H, Riou E, et al. **Toward the definition of cerebral arteriopathies of childhood.** *Curr Opin Pediatr* 2004;16:617–22 [CrossRef Medline](#)
- Sébire G, Meyer L, Chabrier S. **Varicella as a risk factor for cerebral infarction in childhood: a case-control study.** *Ann Neurol* 1999;45:679–80 [CrossRef Medline](#)
- Earley CJ, Kittner SJ, Feeser BR, et al. **Stroke in children and sickle-cell disease: Baltimore-Washington Cooperative Young Stroke Study.** *Neurology* 1998;51:169–76 [CrossRef Medline](#)
- Pandey S, Hakky M, Kwak E, et al. **Application of basic principles of physics to head and neck MR angiography: troubleshooting for artifacts.** *Radiographics* 2013;33:e113–23 [CrossRef Medline](#)
- Corfield L, Speirs A, McCormack DJ, et al. **Time of flight magnetic resonance angiography: a trap for the unwary.** *EJVES Extra* 2010;19:e35–37 [CrossRef](#)
- Takhtani D, Dundamadappa S, Puri AS, et al. **Flow artifact in the anterior communicating artery resembling aneurysm on the time of flight MR angiogram.** *Acta Radiol* 2014;55:1253–57 [CrossRef Medline](#)
- Silvennoinen HM, Ikonen S, Soinne L, et al. **CT angiographic analysis of carotid artery stenosis: comparison of manual assessment, semiautomatic vessel analysis, and digital subtraction angiography.** *AJNR Am J Neuroradiol* 2007;28:97–103 [Medline](#)
- van den Wijngaard IR, Holswilder G, van Walderveen MA, et al. **Treatment and imaging of intracranial atherosclerotic stenosis: current perspectives and future directions.** *Brain Behav* 2016;6:e00536 [CrossRef Medline](#)
- Qureshi AI, Caplan LR. **Intracranial atherosclerosis.** *Lancet* 2014;383:984–98 [CrossRef Medline](#)
- Lan L, Leng X, Abrigo J, et al. **Diminished signal intensities distal to intracranial arterial stenosis on time-of-flight MR angiography might indicate delayed cerebral perfusion.** *Cerebrovasc Dis* 2016;42:232–39 [CrossRef Medline](#)
- Miura M, Nakajima M, Fujimoto A, et al. **Decreased signal intensity ratio on MRA reflects misery perfusion on SPECT in patients with intracranial stenosis.** *J Neuroimaging* 2018;28:206–11 [CrossRef Medline](#)
- Samuels OB, Joseph GJ, Lynn MJ, et al. **A standardized method for measuring intracranial arterial stenosis.** *AJNR Am J Neuroradiol* 2000;21:643–46 [Medline](#)
- Hankey GJ, Warlow CP, Molyneux AJ. **Complications of cerebral angiography for patients with mild carotid territory ischaemia being considered for carotid endarterectomy.** *J Neurol Neurosurg Psychiatry* 1990;53:542–48 [CrossRef Medline](#)

26. Davies KN, Humphrey PR. **Complications of cerebral angiography in patients with symptomatic carotid territory ischaemia screened by carotid ultrasound.** *J Neurol Neurosurg Psychiatry* 1993;56:967–72 [CrossRef Medline](#)
27. Johnston DC, Chapman KM, Goldstein LB. **Low rate of complications of cerebral angiography in routine clinical practice.** *Neurology* 2001;57:2012–14 [CrossRef Medline](#)
28. Liebeskind DS, Kosinski AS, Lynn MJ, et al. **Noninvasive fractional flow on MRA predicts stroke risk of intracranial stenosis.** *J Neuroimaging* 2015;25:87–91 [CrossRef Medline](#)
29. Matsuo S, Nakamizo A, Fujioka Y, et al. **Time-of-flight MRA signal intensity predicts the cerebral hemodynamic status after superficial temporal artery to middle cerebral artery anastomosis.** *J Clin Neurosci* 2019;59:124–29 [CrossRef Medline](#)
30. Leng X, Ip HL, Soo Y, et al. **Interobserver reproducibility of signal intensity ratio on magnetic resonance angiography for hemodynamic impact of intracranial atherosclerosis.** *J Stroke Cerebrovasc Dis* 2013;22:e615–19 [CrossRef Medline](#)
31. Verlhac S, Bernaudin F, Tortrat D, et al. **Detection of cerebrovascular disease in patients with sickle cell disease using transcranial Doppler sonography: correlation with MRI, MRA and catheter angiography.** *Pediatr Radiol* 1995;25:S14–19
32. Kandeel AY, Zimmerman RA, Ohene-Frempong K, et al. **Comparison of magnetic resonance angiography and catheter angiography in sickle cell disease: clinical significance and reliability.** *Neuroradiology* 1996;38:409–16 [CrossRef Medline](#)
33. Koo BK, Erglis A, Doh JH, et al. **Diagnosis of ischemia-causing coronary stenoses by noninvasive fractional flow reserve computed from coronary computed tomographic angiograms: results from the prospective multicenter DISCOVER-FLOW (Diagnosis of Ischemia-Causing Stenoses Obtained Via Noninvasive Fractional Flow Reserve) study.** *J Am Coll Cardiol* 2011;58:1989–97 [CrossRef Medline](#)
34. Min JK, Koo BK, Erglis A, et al. **Usefulness of noninvasive fractional flow reserve computed from coronary computed tomographic angiograms for intermediate stenoses confirmed by quantitative coronary angiography.** *Am J Cardiol* 2012;110:971–76 [CrossRef Medline](#)
35. Gupta A, Chazen JL, Hartman M, et al. **Cerebrovascular reserve and stroke risk in patients with carotid stenosis or occlusion: a systematic review and meta-analysis.** *Stroke* 2012;43:2884–91 [CrossRef Medline](#)
36. Baradaran H, Patel P, Gialdini G, et al. **Quantifying intracranial internal carotid artery stenosis on MR angiography.** *AJNR Am J Neuroradiol* 2017;38:986–90 [CrossRef Medline](#)
37. Chimowitz MI, Lynn MJ, Howlett-Smith H, et al; Warfarin-Aspirin Symptomatic Intracranial Disease Trial Investigators. **Comparison of warfarin and aspirin for symptomatic intracranial arterial stenosis.** *N Engl J Med* 2005;352:1305–16 [CrossRef Medline](#)
38. Hirooka R, Ogasawara K, Inoue T, et al. **Simple assessment of cerebral hemodynamics using single-slab 3D time-of-flight MR angiography in patients with cervical internal carotid artery stenotic diseases: comparison with quantitative perfusion single-photon emission CT.** *AJNR Am J Neuroradiol* 2009;30:559–63 [CrossRef Medline](#)
39. Miralles M, Dolz JL, Cotillas J, et al. **The role of the circle of Willis in carotid occlusion: assessment with phase contrast MR angiography and transcranial duplex.** *Eur J Vasc Endovasc Surg* 1995;10:424–30 [CrossRef Medline](#)