



Discover Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

[VIEW CATALOG](#)

AJNR

**Celebrating 35 Years of the AJNR: May 1986
edition**

AJNR Am J Neuroradiol 2021, 42 (5) 993

doi: <https://doi.org/10.3174/ajnr.P6836>

<http://www.ajnr.org/content/42/5/993>

This information is current as
of September 3, 2025.

Celebrating 35 Years of the AJNR


May 1986 edition

Magnetic Resonance Imaging of Brain Iron

Burton Drayer^{1,2}
 Peter Burger³
 Robert Darwin¹
 Stephen Riederer¹
 Robert Herfkens¹
 G. Allan Johnson¹

A prominently decreased signal in the globus pallidum, red nucleus, and dentate nuclei on T2-weighted images (SE). This MR finding correlated closely with the sites of preferential accumulation of iron in normal postmortem brains. T2 thus provides an accurate in vivo and MR studies in normal brain; the striatum, and still lower level matter, iron concentration is (a) capsule and optic radiations, (b) prominent in the subcortical "ring" in the brain at birth; it increases of brain iron should assist in elucidating neurodegenerative, demyelinating

It has been suggested that exquisite anatomic images, map brain function. Early attempts have been promising but time-consuming using proton MR imaging have but they have been disappointing. When using a high field strength in every patient studied on a finding consisted of decreased specific brain locales, including the globus pallidum, reticular substantia nigra, red nucleus, dentate nucleus, and putamen [1]. This paper gives an account of our studies to determine the origin of this biochemical effect.



This article appears in the May/June 1986 issue of AJNR and the July 1986 issue of AJR. Received October 25, 1985; accepted after revision January 3, 1986.

Presented in part at the annual meeting of the American Society of Neuroradiology, New Orleans, February 1985.

¹ Department of Radiology, Duke University Medical Center, Durham, NC 27710. Address reprint requests to B. P. Drayer, Department of Radiology, Box 3808, Duke University Medical Center, Durham, NC 27710.

² Department of Medicine (Neurology), Duke University Medical Center, Durham, NC 27710.

³ Department of Pathology, Duke University Medical Center, Durham, NC 27710.

AJNR 7:373-380, May/June 1986
 0195-6108/86/0703-0373
 © American Society of Neuroradiology

Magnetic Resonance Imaging of the Chronically Injured Cervical Spinal Cord

Robert M. Quencer¹
 Jerome J. Sheldon²
 M. Judith Donovan Post³
 Rosendo D. Diaz⁴
 Bertie M. Montalvo¹
 Barth A. Green¹
 Frank J. Eismont¹


Thirteen patients evaluated with MRI because of the reliability of our results were compared with in vivo of those the it was found that MRI the injured spinal cord separate myelomalacia frequently difficult to discern with long

Over the years, spinal cord has its percutaneous computed tomography accurate, because media within the spinal cord cyst (2) associated with chronically injured sonography. Spec either myelomalacia therefore, to have patients with shunt but noncystic spinal

Although a number imaging (MRI) in patients not, to our knowledge, injured spinal cords. In this paper we report the use of MRI in 13 patients with prior cervical cord trauma, 10 of whom were also studied with delayed metrizamide CT and five of whom had surgery. It is our objective to compare the results of MRI with those of delayed metrizamide CT in order to determine which is the most accurate method of preoperatively evaluating the chronically damaged spinal cord. Further, we correlate these imaging modalities with the patients' clinical presentations and the findings at surgery as determined by intraoperative spinal sonography.

Materials and Methods

Thirteen patients who had suffered severe cervical spinal cord trauma six months to 11 years prior to their present hospitalization were evaluated clinically and examined radiograph-



This article appears in the May/June 1986 issue of AJNR and the July 1986 issue of AJR. Received August 1, 1985; accepted after revision December 25, 1985.

Presented in part at the annual meeting of the American Society of Neuroradiology, New Orleans, February 1985.

¹ Department of Radiology, University of Miami/Jackson Memorial Medical Center, Miami, FL. Address reprint requests to R. M. Quencer, Department of Radiology (R-130), University of Miami School of Medicine, P.O. Box 016060, Miami, FL 33101.

² Department of Radiology, Mt. Sinai Medical Center, Miami Beach, FL.

³ Department of Neurological Surgery, University of Miami/Jackson Memorial Medical Center, Miami, FL.

⁴ Department of Orthopaedic Surgery, University of Miami/Jackson Memorial Medical Center, Miami, FL.

AJNR 7:457-464, May/June 1986
 0195-6108/86/0704-0457
 © American Society of Neuroradiology