

Providing Choice & Value



Generic CT and MRI Contrast Agents



This information is current as of July 28, 2025.

MR Imaging Findings in Anti-Leucine-Rich Glioma Inactivated Protein 1 Encephalitis: A Systematic Review and Meta-analysis

Francisco C. Almeida, Ana I. Pereira, Catarina Mendes-Pinto, Joana Lopes, João Moura, José Maria Sousa, Gonçalo Videira, Raquel Samões and Tiago Gil Oliveira

AJNR Am J Neuroradiol published online 13 June 2024 http://www.ajnr.org/content/early/2024/06/13/ajnr.A8256 ORIGINAL RESEARCH ULTRA-HIGH-FIELD MRI/IMAGING OF EPILEPSY/DEMYELINATING DISEASES/INFLAMMATION/ INFECTION

MR Imaging Findings in Anti-Leucine-Rich Glioma Inactivated Protein 1 Encephalitis: A Systematic Review and Meta-analysis

Francisco C. Almeida, Ana I. Pereira, Catarina Mendes-Pinto, Joana Lopes, João Moura, José Maria Sousa, Gonçalo Videira,
Raquel Samões, and ¹⁰Tiago Gil Oliveira

ABSTRACT

BACKGROUND: Antibodies against leucine-rich glioma inactivated protein 1 (LGII) constitute a common form of autoimmune encephalitis. On MR imaging, it may show T2 FLAIR hyperintensities of the medial temporal lobe (T2 FLAIR-MTL), involve the basal ganglia, or be unremarkable.

PURPOSE: We performed a systematic review and meta-analysis to obtain prevalence estimates of abnormal findings on MR imaging in anti-LGII encephalitis. A human brain map of the *LGII* microarray gene expression was derived from the Allen Human Brain Atlas.

DATA SOURCES: PubMed and Web of Science were searched with the terms "LGII" and "encephalitis" from inception to April 7, 2022.

STUDY SELECTION: Thirty-one research publications, encompassing case series and retrospective cohort and case-control studies, with >10 patients with anti-LGII encephalitis and MR imaging data were included.

DATA ANALYSIS: Pooled prevalence estimates were calculated using Freeman-Tukey double-arcsine transformation. Meta-analysis used DerSimonian and Laird random effects models.

DATA SYNTHESIS: Of 1318 patients in 30 studies, T2 FLAIR-MTL hyperintensities were present in 54% (95% CI, 0.48–0.60; $I^2 = 76\%$). Of 394 patients in 13 studies, 27% showed bilateral (95% CI, 0.19–0.36; $I^2 = 71\%$) and 24% unilateral T2 FLAIR-MTL abnormalities (95% CI, 0.17–0.32; $I^2 = 61\%$). Of 612 patients in 15 studies, basal ganglia abnormalities were present in 10% (95% CI, 0.06–0.15; $I^2 = 67\%$). *LGII* expression was highest in the amygdala, hippocampus, and caudate nucleus.

LIMITATIONS: Only part of the spectrum of MR imaging abnormalities in anti-LGII encephalitis could be included in a meta-analysis. MR imaging findings were not the main outcomes in most studies, limiting available information. I^2 values ranged from 62% to 76%, representing moderate-to-large heterogeneity.

CONCLUSIONS: T2 FLAIR-MTL hyperintensities were present in around one-half of patients with anti-LGII. The prevalence of unilateral and bilateral presentations was similar, suggesting unilaterality should raise the suspicion of this disease in the appropriate clinical context. Around 10% of patients showed basal ganglia abnormalities, indicating that special attention should be given to this region. *LGII* regional expression coincided with the most frequently reported abnormal findings on MR imaging. Regional specificity might be partially determined by expression levels of the target protein.

ABBREVIATIONS: ADAM = a disintegrin and metalloprotease domain; AIE = autoimmune encephalitis; AMPA = α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; BG = basal ganglia; CASPR2 = contactin-associated protein-like 2; EEG = electroencephalogram; FBDS = faciobrachial dystonic seizures; LGII = leucine-rich glioma inactivated protein 1; MTL = medial temporal lobe

A ntibodies against leucine-rich glioma inactivated protein 1 (LGI1) characterize the second most common form of

Received November 14, 2023; accepted after revision February 14, 2024. From the Departments of Neuroradiology (F.C.A., A.I.P., C.M.-P.) and Neurology (J.L., J.M., G.V., R.S.), Centro Hospitalar Universitàrio de Santo António, Porto, Portugal; Life and Health Science Research Institute (F.C.A., T.G.O.), School of Medicine, University of Minho, Braga, Portugal; Life and Health Sciences Research Institute/38's-PT Government Associate Laboratory (F.C.A., T.G.O.), Braga/Guimarães, Portugal; Department of Neuroradiology (J.M.S.), Centro Hospitalar Universitário de São João, Porto, Portugal; Unit for Multidisciplinary Research in Biomedicine (R.S.), Instituto de Ciências Biomédicas de Abel Salazar da Universidade do Porto, Porto, Portugal; and Department of Neuroradiology (T.G.O.), Hospital de Braga, Braga, Portugal. autoimmune encephalitis (AIE).¹ These antibodies were first described in 2010 and were previously clustered within voltagegated potassium channel antibodies, which included contactinassociated protein-like 2 (CASPR2) along with LGI1.² The LGI1

Raquel Samões and Tiago Gil Oliveira contributed equally to this work.

Please address correspondence to Francisco C. Almeida, MD, Neuroradiology Department, Largo do Prof. Abel Salazar, 4099-001 Porto, Portugal; e-mail: franciscoalmeida.neurorradiologia@chporto.min-saude.pt

Indicates article with online supplemental data.

http://dx.doi.org/10.3174/ajnr.A8256

AJNR Am J Neuroradiol •:• • 2024 www.ajnr.org 1



FIG 1. Illustrative examples of signal abnormalities in 2 cases of confirmed anti-LGII encephalitis, one with medial temporal lobe T2-FLAIR hyperintensities (*A*) and another with right BG T2-FLAIR hyperintensity (*B*).

protein is widely expressed in the CNS and particularly in neurons, where it forms a trans-synaptic protein complex that regulates ion channel and synaptic activity.^{3,4}

This disease typically presents with subacute cognitive impairment and epilepsy, but faciobrachial dystonic seizures (FBDS) are the most distinctive clinical feature. Although presenting across a wide variety of ages, it is thought to be most common in those older than 60 years of age.³ The clinical presentation thus leads to a vast differential diagnosis, in which clinical suspicion plays an important role.^{5,6} MR imaging is relevant in supporting the diagnosis of AIE and excluding alternative diagnoses. Characteristically, it involves medial temporal lobe (MTL) hyperintensities on T2 FLAIR sequences, which represent a criterion for a definite diagnosis of AIE when observed bilaterally (Fig 1A).^{7,8} This laterality issue is because in the absence of antibody confirmation, unilateral hippocampal/MTL abnormalities have a wider radiologic differential diagnosis.^{7,9} The main diagnostic considerations in this imaging presentation include seizure-related MR imaging abnormalities;¹⁰ viral encephalitis, particularly herpes simplex virus 1 and 2 and human herpesvirus 6;¹¹ neurosyphilis;¹² Whipple disease,¹³ temporal lobe glioma,¹⁴ and acute posterior circulation ischemic infarction.¹⁵

However, not all cases of anti-LGI1 encephalitis present with MTL abnormalities at the time of MR imaging. MR imaging can also be unremarkable or show basal ganglia (BG) involvement (Fig 1*B*).¹⁶ Moreover, given the relatively recent description of the wide range of antibodies causing AIE, MR imaging findings are not yet thoroughly described for each antibody group.^{8,17} It is, therefore, important to provide prevalence estimates of MR imaging abnormalities in the literature for each type of antibody-mediated encephalitis. Indeed, given the wide clinical and brain imaging differential diagnoses involved in these cases, detailing the prevalence of specific patterns of abnormal findings on MR imaging for each encephalitis might aid the radiologist in providing a narrower differential while waiting for laboratory results.

Furthermore, abnormalities on MR imaging are likely reflective of pathophysiologic mechanisms. Characterizing the topography of the most frequently reported abnormalities may shed light on the underlying causative pathology. Although most studies have focused on *LGI1* expression in rodent models derived from immunohistochemistry or patient-derived antibodies,^{3,4} a human brain map of the *LGI1* gene might reveal why selective brain areas are impacted in this disease.

In this study, we performed a systematic review and meta-analysis of the proportion of MTL and BG MR imaging signal abnormalities in LGI1 encephalitis. Moreover, we used the publicly accessible Allen Human Brain Atlas (https://human.brain-map.org) to derive a *LGI1* gene expression map

of the human brain, which can provide potential hints as to why LGI1 encephalitis preferentially affects certain brain regions.

MATERIALS AND METHODS

This systematic review and meta-analysis was conducted in accordance with the PRISMA guidelines. 18

Search Strategy and Selection Criteria

We searched PubMed and the Web of Science from inception to April 7, 2022, using the search terms "LGI1" and "encephalitis." PubMed had 511 publications and the Web of Science had 646 publications. Bibliographic sources from selected articles were also used to assess research publications for eligibility. Original research articles, including case series, retrospective or prospective cohort studies, and case-control studies were considered for selection. Case reports, systematic reviews, meta-analyses, review articles, and other nonoriginal research publications were excluded during screening. Only studies conducted with human participants and published in English were considered. The number of each article type found during screening is provided in the Online Supplemental Data.

Inclusion and Exclusion Criteria

Research publications were eligible if they included a clinical population of LGI1 encephalitis and reported >10 cases, when revealed in the abstract. The clinical definition of LGI1 encephalitis was then confirmed on inspection of the publications to comply with definite autoimmune encephalitis, requiring detection of anti-LGI1 antibodies on CSF and/or serum.⁷ Exclusion criteria after retrieval were the following: the study did not present MR imaging data, did not clearly specify the MR imaging abnormalities/brain location/MR imaging sequence used, did not use T2-FLAIR sequences or used other types of MR imaging data not widely used in clinical practice, had \leq 10 patients with LGI1 encephalitis, did not separate LGI1 from CASPR2 encephalitis or from other autoantibodies; and the data were previously published in other research publications, there was no accessibility of the full text, or the article had low manuscript quality.

Data Extraction

Two authors independently screened titles and abstracts and assessed the full text for eligibility (F.C.A., A.I.P). Discrepancies were solved after discussion between the 2 authors. Clinical data (J.L., J.M.), electroencephalogram (EEG) data (G.V., F.C.A.), and neuroimaging data (F.C.A., C.M.-P.) were extracted independently by 2 authors. When necessary, discrepancies were solved by a third party (clinical: R.S.; neuroimaging: T.G.O.). The quality of the studies was assessed independently by 2 authors (F.C.A. and A.I.P.) on the basis of a previously published tool for prevalence studies.¹⁹ This tool consists of 10 items consisting of 4 external validity and 6 internal validity items, the sum score of which provides a summary assessment of the risk of bias.

Data Analysis

Pooled prevalence estimates of neuroimaging findings were calculated using Freeman-Tukey double-arcsine transformation. Metaanalysis was conducted using DerSimonian and Laird randomeffects models. Heterogeneity was calculated using I² statistics, and the Cochran Q test was used to assess statistical significance. The metaprop package in R (https://www.rdocumentation.org/ packages/meta/versions/6.2-1/topics/metaprop) was used for this analysis.

Metaregression analysis was conducted to assess modifiers of prevalence. Clinical symptoms and CSF antibody positivity were used in univariate models to assess the effects on the prevalence of both T2 MTL hyperintensities and BG signal abnormalities (T2 hyperintensities and T1 hypo- and hyperintensities). Studies with missing data on modifier variables were excluded. R^2 was used to assess the proportion of the heterogeneity explained by the modifier, the *P* value of the Quantitative Methods statistics was used for statistical significance, and I², for residual heterogeneity left unexplained. The Meta package in R was used for this analysis (https://cran.r-project.org/web/packages/meta/index.html).

Publication bias was assessed with visual inspection of funnel plots and assessed for statistical significance using the Egger weighted linear regression test. The meta package in R was used for this analysis.

LGI1 Gene Brain-Expression Map

Regional microarray expression data were obtained from 6 postmortem brains (1 woman; age range, 24.0–57.0 years; mean age, 42.50 [SD, 13.38] years) provided by the Allen Human Brain Atlas.²⁰ Data were processed with the abagen toolbox (version 0.1.3; https://github.com/rmarkello/abagen) using an 83-region volumetric atlas in Montreal Neurological Institute space. First, microarray probes were reannotated using data provided by Arnatkeviciute et al.²¹ Probes not matched to a valid Entrez ID were discarded. Next, probes were filtered on the basis of their expression intensity relative to background noise²² so that probes with intensity less than the background in \geq 50% of samples across donors were discarded, yielding 31,569 probes. When multiple probes indexed the expression of the same gene, we selected and used the probe with the most consistent pattern of regional variation across donors (ie, differential stability²³). Here, regions correspond to the structural designations provided in the ontology from the Allen Human Brain Atlas. The Montreal Neurological Institute coordinates of tissue samples were updated to those generated via nonlinear registration using the Advanced Normalization Tools (ANTs; http://stnava.github. io/ANTs/). Samples were assigned to brain regions in the provided atlas if their Montreal Neurological Institute coordinates were within 2 mm of a given parcel.

To reduce the potential for misassignment, sample-to-region matching was constrained by hemisphere and gross structural divisions (ie, cortex, subcortex/brainstem, and cerebellum, such that for example, a sample in the left cortex could only be assigned to an atlas parcel in the left cortex²¹). All tissue samples not assigned to a brain region in the provided atlas were discarded. Intersubject variation was addressed by normalizing tissue sample expression values across genes using a robust sigmoid function.²⁴ Normalized expression values were then rescaled to the unit interval. Gene expression values were then normalized across tissue samples using an identical procedure. Samples assigned to the same brain region were averaged separately for each donor and then across donors, yielding a regional expression matrix with 83 rows corresponding to brain regions and 15,633 columns corresponding to the retained genes. Normalized gene expression values were then plotted in a brain map using the ggseg R package (https://cran.r-project.org/web/packages/ ggseg/index.html)²⁵ with a Desikan-Killiany atlas.

Data Availability

Data used in this meta-analysis will be provided at request.

RESULTS

Study and Clinical Characteristics

In this study, we included 31 research publications (Fig 2).^{5,16,26-54} Two studies reported partially on the same patients but focused on different neuroimaging findings.^{16,50} For summary statistics regarding each of the clinical and neuroimaging findings, we excluded the publication with fewer patients. In total, the sample was composed of 1409 patients with anti-LGI1 encephalitis. The mean quality-assessment score of the research publications was 3, indicating overall good quality (Online Supplemental Data).

Most of the studies were retrospective and based on clinical cohorts, with some studies drawing their sample from EEG or PET cohorts. Case definition was mostly based on criteria for definite AIE,⁷ with 3 studies basing their definition on FBDS and LGI1 antibody positivity.^{16,43,49}

Table 1 reports summary statistics for the whole study population. The mean age was 61.4 years, and 64% of patients were men. Seventy-eight percent of the patients presented with cognitive impairment (1095 of 1409). For FBDS, 27 studies presented data, totaling 1252 patients, among whom 639 (51%) had FBDS at any point in time. Twenty-eight studies reported seizures other than FBDS, totaling 1368 patients, of whom 989 (72%) were affected. Twenty-four studies, encompassing 1146 patients included data on hyponatremia, among whom 512 (45%) patients presented with low serum sodium. Consciousness impairment was evaluated in 12 studies with 574 patients and was present in



FIG 2. Study selection flow chart.¹⁸ PubMed and Web of Science were searched for the terms "LGII" and "encephalitis." Of 665 records screened, 133 were assessed for eligibility, resulting in 31 included publications.

Table 1: Summary statistics for the whole study population

	Studies (No.)	Total Subjects (No.)	Summary Statistic
Mean age (yr)	30	1409	61.4
Male sex (No.) (%)	30	1409	895 (64%)
Cognitive impairment (No.) (%)	30	1409	1095 (78%)
FBDS (No.) (%)	27	1252	639 (51%)
Other seizures (No.) (%)	28	1368	989 (72%)
Psychiatric symptoms (No.) (%)	25	1203	612 (51%)
Hallucinations (No.) (%)	9	493	80 (16%)
Hyponatremia (No.) (%)	24	1146	512 (45%)
Sleep disorder (No.) (%)	21	1166	417 (36%)
Consciousness impairment (No.) (%)	12	574	180 (31%)
Autonomic dysfunction (No.) (%)	15	812	155 (19%)
Abnormal EEG findings (No.) (%)	19	698	508 (73%)
CSF antibody positivity (No.) (%)	14	713	457 (64%)

180 (31%). Autonomic dysfunction was present in 155 patients of 812 (19%) in 15 studies. Sleep disorder was present in 417 patients of 1166 (36%) in the 21 studies that reported data on this symptom. Of 1203 patients, 612 in 25 studies presented with psychiatric symptoms (51%). Hallucinations were present in 80 of 493 patients (16%) in 9 studies. Personality changes and psychosis (unspecified) were among other symptoms reported in the included literature. In 14 studies with 713 patients, CSF antibody positivity was found in 457 (64%). In 19 studies with data on EEG, 508 of 698 patients (73%) presented with abnormal findings.

Neuroimaging Findings

Of the whole sample of 1409 patients, 1318 had reported data on MR imaging. Thirty studies reported data on T2 or T2 FLAIR-MTL hyperintensities. Meta-analysis of the proportion of patients with T2 FLAIR-MTL revealed that this finding was present in 54% of patients (Fig 3; 95% CI, 0.48–0.60; $I^2 = 76\%$). Thirteen

studies reported T2 FLAIR-MTL laterality with a total of 394 patients. Metaanalysis revealed 27% with bilateral T2-MTL findings (Fig 4*A*; 95% CI, 0.19–0.36; $I^2 = 71\%$) and 24% with unilateral abnormalities (Fig 4*B*; 95% CI, 0.17–0.32; $I^2 = 61\%$) of the whole population (ie, with and without T2-MTL abnormalities).

Fifteen studies encompassing 612 patients reported data on BG MR imaging signal abnormalities, which included T2 hyperintensities and T1 hypo- and hyperintensities. Altogether, meta-analysis of proportions revealed these signal abnormalities to be present in 10% of patients (Fig 5; 95% CI, 0.06–0.15; $I^2 = 67\%$).

Inspection of funnel plots and linear regression testing for funnel asymmetry did not reveal a significant potential for publication bias (Online Supplemental Data).

Although not reported frequently enough to undergo meta-analysis, studies showed anti-LGI1 encephalitis with other regional and MR abnormalities.^{26-38,40,41,43,45-48,51,54} Amygdala, insula, and other neocortical T2-FLAIR hyperintensities were found in around 10%, 8%, and 18% of patients, respectively (Table 2). Two patients presented with cortical diffusion restriction (2%). Leptomeningeal gadolinium enhancement was described in 3 patients (5%). MTL swelling was reported in 18 patients (15%), and atrophy, in 55 patients (16%), whereas gadolinium enhancement was found in 5 patients (3%). In the studies with available

longitudinal follow-up, 46% of patients of 154 in 6 studies developed atrophy or mesial temporal sclerosis. BG atrophy was reported in 3 patients (3%). Neocortical and cerebellum gadolinium enhancement was described in 1 patient each in 1 study.²⁷ Diffusion restriction in the basal ganglia was reported in 1 study.¹⁶

Given the high heterogeneity found in the meta-analysis of these neuroimaging findings, we performed meta-regression analysis for T2-MTL and BG signal abnormalities with clinical predictors (Online Supplemental Data). We did not find any statistically significant modifiers of the proportion of MR imaging signal abnormalities.

LGI1 Gene Expression Map of the Human Brain

To assess whether the *LGI1* gene is differentially expressed in the selective regions affected on MR imaging, we next used publicly available data from the Allen Human Brain Atlas to create a

Medial Temporal Lobe T2-FLAIR Hyperintensities



FIG 3. Forest plot showing the prevalence of T2 FLAIR-MTL hyperintensities. Of 1318 patients in 30 studies, 54% presented with this signal abnormality.

gene-expression brain map. We found that *LGI1* is expressed widely across the cortex but is particularly predominant in subcortical regions, namely, the amygdala, hippocampus, and caudate nucleus (Fig 6*A*,-*B*; and Online Supplemental Data).

DISCUSSION

In this systematic review and meta-analysis, we focused on neuroimaging findings of anti-LGI1 encephalitis. We included 31 studies and 1409 patients. Our studied population had a mean age of 61.4 years and a mild predominance of male patients (64%). The most frequently reported symptoms were cognitive impairment, seizures other than FBDS, followed by FBDS and psychiatric symptoms. These clinical and demographic characteristics are in agreement with those in previous studies.⁵⁵

We found that 54% of patients presented with T2-FLAIR hyperintensities of the MTL (Fig 3). This finding reinforces the idea that a negative MR imaging in anti-LGI1 encephalitis is frequent. In a subgroup analysis, unilateral and bilateral presentations were similar, with a prevalence of 24% and 27%, respectively (Fig 4*A*, -*B*). This finding suggests that in the appropriate clinical context, the presence of unilateral MTL abnormalities on MR imaging should raise the possibility of AIE, namely anti-LGI1 encephalitis, and may aid in therapeutic decisions while awaiting antibody confirmation. A common imaging differential diagnosis for T2 FLAIR-MTL hyperintensities frequently associated with volume expansion includes other autoimmune/paraneoplastic

encephalitis cases, among which the neuronal-surface antigen group includes NMDAR,⁵⁶ CASPR2,^{57,58} GABA-B,^{59,60} Neurexin-3a,⁶¹ AMPAR,^{62,63} mGluR5,⁶⁴ GlyR,65,66 AK5,67 and DPPX6,68 and in the intracellular antigen group includes GAD65,⁶⁹ Hu,⁸ amphiphysin,⁷⁰ and Ma2.⁷¹ Viral encephalitis,¹¹ neurosyphilis,12 Whipple disease,13 glioma,14 ictal changes,9 and posterior circulation stroke¹⁵ are among other important diagnostic considerations. Depending on the time of imaging, anti-LGI1 encephalitis might already have presented with hippocampal atrophy, in which case neurodegenerative diseases, including Alzheimer disease,^{72,73} frontotemporal dementia,⁷⁴ and limbic-predominant age-related TDP-43 encephalopathy,75 as well as medial temporal sclerosis of other causes, are to be included in the differential diagnosis. Other causes to consider, depending on clinical presentation and other associated signal changes, include metabolic and mitochondrial disorders.⁷⁶

Notably, BG signal abnormalities, encompassing both T2 and T1 hyperintensities and T1 hypointensities, were reported in around 10% of

patients (Fig 5). The available number of studies providing information regarding the BG was substantially smaller than that for the MTL. However, these changes can be the sole signal alteration observed on MR imaging and might be unilateral¹⁶ (Fig 1*B*), suggesting that this brain region deserves detailed attention when suspecting AIE. In fact, LGI1 expression is high in the BG in humans (Fig 6*B*), and [¹⁸F] fluorodeoxyglucose PET studies have repeatedly shown metabolic alterations in these regions.^{27,39,41} Most important, these lesions can present with diffusion restriction,¹⁶ the cause of which is unknown. Ischemic infarction is thus an important differential diagnosis, and attention to clinical presentation is crucial.

The nature of T1 hyperintensities of the BG is also not yet understood. Underlying causes of T1 relaxation time shortening included blood, fat, high paramagnetic substances such as iron, and high protein content. In anti-LGI1 encephalitis, BG T1hyperintensities are sometimes preceded by T2-hyperintensities and have been shown to remain for a longer time than the latter before resolving.¹⁶ A putative explanation might be the accumulation of proteins at the site of antibody engagement, which persists after initial inflammatory changes. A parallel explanation has been drawn for T1 pallidal hyperintensities in Creutzfeldt-Jakob disease, where accumulation of prion protein, confirmed at neuropathology, was correlated with this MR imaging signal abnormality.⁷⁷ Signal alterations in the BG have a vast differential diagnosis, which includes toxic and metabolic disorders,⁷⁸ among which hepatic failure and Wilson disease are important



Unilateral Medial Temporal Lobe T2-FLAIR Hyperintensities



FIG 4. Forest plots showing the prevalence of bilateral T2 FLAIR-MTL hyperintensities (27% of 394 patients in 13 studies) (*A*) and unilateral MTL T2 FLAIR-MTL hyperintensities (24% of 394 patients in 13 studies) (*B*).



FIG 5. Forest plot showing the prevalence of BG signal abnormalities (T2-FLAIR hyperintensities and T1 hypo- and hyperintensities with 10% of 612 patients having these abnormalities).

considerations, Creutzfledt-Jakob disease,⁷⁹ other autoimmune encephalitis (eg, CV2/collapsin response mediated protein 5),⁸⁰ viral encephalitis,¹¹ and arterial and venous infarction, among others.⁸¹

aptic activity.³ Evidence from animal models has shown that patient-derived anti-LGI1 antibodies induce important impairment of synaptic plasticity,^{83,84} neuronal excitability⁸⁵ and memory in rodents.^{83,84} This evidence supports the pathogenicity of the

Other brain regions were found to be altered in LGI1 encephalitis, namely, the amygdala, insula, cerebellum, and different regions of the neocortex, with signal abnormalities across T1 with gadolinium enhancement, T2 and diffusion MR imaging, as well as leptomeningeal gadolinium enhancement. This finding suggests that LGI1 AIE can present with a widespread variety of signal abnormalities across multiple brain regions and multiple MR imaging sequences. Moreover, the under-reporting of altered signal beyond the MTL indicates that there is a need for further research in systematizing MR imaging signal abnormalities in LGI1 encepha-

In the studies that included a follow-up period, 46% of patients went on to develop hippocampal atrophy or mesial temporal sclerosis (Table 2). These results support the concept that anti-LGI1 encephalitis might induce a neurodegenerative process in a large proportion of patients, which might be associated with remaining cognitive deficits.⁸² Detection of these imaging findings by the radiologist in serial scans provides useful information for the clinician.

litis across the whole brain.

To assess qualitatively whether MR imaging abnormalities tend to occur at brain regions most enriched for LGI1 gene expression, we generated a brain map using the publicly available Allen Human Brain Atlas. Regions with the highest expression were the amygdala, hippocampus, and caudate (Fig 6), suggesting that at least partially, the selective appearance of abnormalities on MR imaging might be due to the relative expression of LGI1. The LGI1 protein is expressed at the synapse, where binding presynaptic a disintegrin and metalloprotease domain 23 (ADAM23) and postsynaptic ADAM22 forms a trans-synaptic tether that couples presynaptic voltage-gated potassium channels and postsynaptic AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) receptors, thereby regulating syn-

Studies (No.)	Total Subjects (No.)	Proportion
3	94	9 (10%)
5	153	12 (8%)
4	79	14 (18%)
3	85	2 (2%)
2	61	3 (5%)
4	117	18 (15%)
7	353	55 (16%)
2	151	5 (3%)
6	154	71 (46%)
2	102	3 (3%)
	Studies (No.) 3 5 4 3 2 4 7 2 6 2	Studies (No.) Total Subjects (No.) 3 94 5 153 4 79 3 85 2 61 7 353 2 117 7 353 2 151 6 154 2 102

^a Excluding MTL and insula.

^b The time of the MR imaging relative to symptom initiation was not available



FIG 6. *LGI*¹ gene expression in the cortical (*A*) and subcortical (*B*) regions derived from the Allen Human Brain Atlas. In this figure, *LGI*¹ gene expression is shown to be higher in the MTLs and BG, concordant with the main MR imaging signal abnormality findings, suggesting that regional brain susceptibility in anti-LGI1 encephalitis might be mediated by the relative levels of brain regional expression of this gene.

antibody and the hypothesis that higher expression of the target antigen might be a contributing factor to explain brain regional susceptibility in these diseases. The accessibility of the antibodies to these brain regions is also a putative contributing factor. Many patients presenting without MR imaging signal abnormalities suggests that only in certain cases is the underlying pathophysiologic mechanism capable of generating a detectable change on MR imaging. The reasons are still to be explored. PET imaging approaches might unravel promising biologic markers of disease activity in the acute phase.⁸⁶

While these results provide the first estimate of neuroimaging signal abnormalities reported in the literature for LGI1 encephalitis, this study has important limitations. Even though MR imaging signal abnormality is likely to be present at the time of symptom onset, we were unable to report any analysis on this parameter. Moreover, neuroimaging findings were not the main outcome in most studies. While we did not find any clinical modifiers on the prevalence of these neuroimaging findings, not all studies provided data for all subjects who underwent MR imaging. More research is needed to establish whether signal abnormalities on MR imaging correlate with clinical presentation or other auxiliary tests. Finally, the specific MR imaging sequence parameters and magnetic field strength of the scanner were not taken into account in the analyses.

Most important, a considerable number of studies were excluded due to incomplete reporting on MR imaging data (Fig 2). These results lead us to highlight the need to include detailed MR imaging descriptions in a disease that can present with a variety of signal abnormalities across various MR imaging sequences.

CONCLUSIONS

This study presents a first approach to estimate the main findings of anti-LGI1 AIE on MR imaging and underlines the importance of considering this diagnosis in the context of unilateral signal abnormalities of the MTL, while also calling for detailed attention to the BG with suspicion of this disease. Most interesting, these regions

express the *LGI1* gene at high levels, thus lending support to the hypothesis that the regions affected by autoimmunity might be at least partially determined by the expression levels of the target protein.

ACKNOWLEDGMENTS

We thank Dr Alexandra Rodrigues from the Neuroradiology Department of Centro Hospitalar Universitário de São José for her valuable input.

 ${\sf Disclosure\ forms\ provided\ by\ the\ authors\ are\ available\ with\ the\ full\ text\ and\ PDF\ of\ this\ article\ at\ www.ajnr.org.$

REFERENCES

- 1. Hébert J, Riche B, Vogrig A, et al. Epidemiology of paraneoplastic neurologic syndromes and autoimmune encephalitides in France. *Neurol Neuroimmunol Neuroinflamm* 2020;7:e883 CrossRef Medline
- Irani SR, Alexander S, Waters P, et al. Antibodies to Kv1 potassium channel-complex proteins leucine-rich, glioma inactivated 1 protein and contactin-associated protein-2 in limbic encephalitis, Morvan's syndrome and acquired neuromyotonia. *Brain* 2010;133:2734–48 CrossRef Medline
- Baudin P, Cousyn L, Navarro V. The LGI1 protein: molecular structure, physiological functions and disruption-related seizures. *Cell Mol Life Sci* 2021;79:16 CrossRef Medline
- Fels E, Muñiz-Castrillo S, Vogrig A, et al. Role of LG11 protein in synaptic transmission: from physiology to pathology. *Neurobiol Dis* 2021;160:105537 CrossRef Medline
- Bastiaansen AE, van Steenhoven RW, de Bruijn MA, et al. Autoimmune encephalitis resembling dementia syndromes. Neurol Neuroimmunol Neuroinflamm 2021;8:e1039 CrossRef Medline
- Costa D, Sardoeira A, Carneiro P, et al. Autoimmune encephalitis: suspicion in clinical practice and mimics. J Neuroimmunol 2022;365:577824 CrossRef Medline
- Graus F, Titulaer MJ, Balu R, et al. A clinical approach to diagnosis of autoimmune encephalitis. *Lancet Neurol* 2016;15:391–404 CrossRef Medline
- 8. Kelley BP, Patel SC, Marin HL, et al. Autoimmune encephalitis: pathophysiology and imaging review of an overlooked diagnosis. *AJNR Am J Neuroradiol* 2017;38:1070–78 CrossRef Medline
- 9. Alves IS, Coutinho AM, Vieira AP, et al. **Imaging aspects of the hippocampus.** *Radiographics* 2022;42:822–40 CrossRef Medline
- Kim SE, Lee BI, Shin KJ, et al. Characteristics of seizure-induced signal changes on MRI in patients with first seizures. Seizure 2017;48:62–68 CrossRef Medline
- 11. Ramli NM, Bae YJ. Structured imaging approach for viral encephalitis. *Neuroimaging Clin N Am* 2023;33:43–56 CrossRef Medline
- Abdelerahman KT, Santamaria DD, Rakocevic G. Pearls and oysters: neurosyphilis presenting as mesial temporal encephalitis. *Neurology* 2012;79:e206–08 CrossRef Medline
- Black DF, Aksamit AJ, Morris JM. MR imaging of central nervous system Whipple disease: a 15-year review. *AJNR Am J Neuroradiol* 2010;31:1493–97 CrossRef Medline
- Zoccarato M, Valeggia S, Zuliani L, et al. Conventional brain MRI features distinguishing limbic encephalitis from mesial temporal glioma. *Neuroradiology* 2019;61:853–60 CrossRef Medline
- Salerno A, Strambo D, Nannoni S, et al. Patterns of ischemic posterior circulation strokes: aclinical, anatomical, and radiological review. *Int J Stroke* 2022;17:714–22 CrossRef Medline
- Flanagan EP, Kotsenas AL, Britton JW, et al. Basal ganglia T1 hyperintensity in LGI1-autoantibody faciobrachial dystonic seizures. Neurol Neuroinflamm 2015;2:e161 CrossRef Medline
- Dalmau J, Graus F. Antibody-mediated encephalitis. N Engl J Med 2018;378:840–51 CrossRef Medline
- Moher D, Liberati A, Tetzlaff J, et al; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009;6:e1000097 CrossRef Medline
- Hoy D, Brooks P, Woolf A, et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. J Clin Epidemiol 2012;65:934–39 CrossRef Medline

- 20. Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, et al. An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* 2012;489:391–99 CrossRef Medline
- Arnatkeviciute A, Fulcher BD, Fornito A. A practical guide to linking brain-wide gene expression and neuroimaging data. *NeuroImage* 2019;189:353–67 CrossRef Medline
- Quackenbush J. Microarray data normalization and transformation. Nat Genet 2002;32(Suppl):496–501 CrossRef Medline
- 23. Hawrylycz M, Miller JA, Menon V, et al. Canonical genetic signatures of the adult human brain. Nat Neurosci 2015;18:1832–44 CrossRef Medline
- 24. Fulcher BD, Little MA, Jones NS. Highly comparative time-series analysis: the empirical structure of time series and their methods. J R Soc Interface 2013;10:20130048 CrossRef Medline
- 25. Mowinckel AM, Vidal-Piñeiro D. Visualization of brain statistics with R packages ggseg and ggseg3d. Advances in Methods and Practices in Psychological Science 2020;3:466–83 CrossRef
- 26. Guo K, Liu X, Lin J, et al. Clinical characteristics, long-term functional outcomes and relapse of anti-LGI1/Caspr2 encephalitis: a prospective cohort study in Western China. Ther Adv Neurol Disord 2022;15:17562864211073203 CrossRef Medline
- 27. Rissanen E, Carter K, Cicero S, et al. **Cortical and subcortical dysmetabolism are dynamic markers of clinical disability and course in anti-LGI1 encephalitis.** *Neurol Neuroimunol Neuroinflamm* 2022;9: e1136 CrossRef Medline
- Yang M, Lian Y. Clinical features and early recognition of 242 cases of autoimmune encephalitis. *Front Neurol* 2021;12:803752 CrossRef Medline
- Cui L, Boltze J, Zhang Y. Positive LG11 antibodies in CSF and relapse relate to worse outcome in anti-LG11 encephalitis. *Front Immunol* 2021;12:772096 CrossRef Medline
- Nissen MS, Ryding M, Nilsson AC, et al. CSF-neurofilament light chain levels in NMDAR and LGI1 encephalitis: a national cohort study. Front Immunol 2021;12:719432 CrossRef Medline
- 31. Zhao X, Zhao S, Chen Y, et al. Subcortical hypermetabolism associated with cortical hypometabolism is a common metabolic pattern in patients with anti-leucine-rich glioma-inactivated 1 antibody encephalitis. Front Immunol 2021;12:672846 CrossRef Medline
- 32. Chen W, Wang M, Gao L, et al. Neurofunctional outcomes in patients with anti-leucine-rich glioma inactivated 1 encephalitis. *Acta Neurol Scand* 2021;144:632–39 CrossRef Medline
- 33. Lin N, Liu Q, Chen J, et al. Long-term seizure outcomes in patients with anti-leucine-rich glioma-inactivated 1 encephalitis. *Epilepsy Behav* 2021;122:108159 CrossRef Medline
- 34. Li TR, Zhang YD, Wang Q, et al. Clinical characteristics and longterm prognosis of anti-lgi1 encephalitis: a single-center cohort study in Beijing, China. Front Neurol 2021;12:674368 CrossRef Medline
- 35. Lardeux P, Fourier A, Peter E, et al. Core cerebrospinal fluid biomarker profile in anti-LGI1 encephalitis. J Neurol 2022;269:377–88 CrossRef Medline
- Muñiz-Castrillo S, Haesebaert J, Thomas L, et al. Clinical and prognostic value of immunogenetic characteristics in anti-LGI1 encephalitis. Neurol Neuroimmunol Neuroinflamm 2021;8:e974 CrossRef Medline
- Zhao Q, Sun L, Zhao D, et al. Clinical features of anti-leucine-rich glioma-inactivated 1 encephalitis in northeast China. Clin Neurol Neurosurg 2021;203:106542 CrossRef Medline
- 38. Shao X, Fan S, Luo H, et al. Brain magnetic resonance imaging characteristics of anti-leucine-rich glioma-inactivated 1 encephalitis and their clinical relevance: a single-center study in China. *Front Neurol* 2020;11:618109 CrossRef Medline
- 39. Liu X, Shan W, Zhao X, et al. The clinical value of ¹⁸F-FDG-PET in autoimmune encephalitis associated with LGI1 antibody. Front Neurol 2020;11:418 CrossRef Medline
- 40. Liu X, Han Y, Yang L, et al. The exploration of the spectrum of motor manifestations of anti-LG11 encephalitis beyond FBDS. *Seizure* 2020;76:22–27 CrossRef Medline

- 41. Jang Y, Lee ST, Bae JY, et al. LGI1 expression and human brain asymmetry: insights from patients with LGI1-antibody encephalitis. J Neuroinflammation 2018;15:279 CrossRef Medline
- 42. Feyissa AM, Lamb C, Pittock SJ, et al. Antiepileptic drug therapy in autoimmune epilepsy associated with antibodies targeting the leucine-rich glioma-inactivated protein 1. *Epilepsia Open* 2018;3:348– 56 CrossRef Medline
- 43. Thompson J, Bi M, Murchison AG, et al; Faciobrachial Dystonic Seizures Study Group. The importance of early immunotherapy in patients with faciobrachial dystonic seizures. *Brain* 2017;141:348– 56 CrossRef Medline
- 44. Celicanin M, Blaabjerg M, Maersk-Moller C, et al. Autoimmune encephalitis associated with voltage-gated potassium channels-complex and leucine-rich glioma-inactivated 1 antibodies: a national cohort study. Eur J Neurol 2017;24:999–1005 CrossRef Medline
- 45. Kim TJ, Lee ST, Moon J, et al. Anti-LGI1 encephalitis is associated with unique HLA subtypes. Ann Neurol 2017;81:183–92 CrossRef Medline
- 46. Finke C, Prüss H, Heine J, et al. Evaluation of cognitive deficits and structural hippocampal damage in encephalitis with leucine-rich, glioma-inactivated 1 antibodies. JAMA Neurol 2017;74:50–59 CrossRef Medline
- van Sonderen A, Thijs RD, Coenders EC, et al. Anti-LGI1 encephalitis: clinical syndrome and long-term follow-up. *Neurology* 2016;87:1449–56 CrossRef Medline
- Navarro V, Kas A, Apartis E, et al; collaborators. Motor cortex and hippocampus are the two main cortical targets in LGI1-antibody encephalitis. Brain 2016;139:1079–93 CrossRef Medline
- Irani SR, Michell AW, Lang B, et al. Faciobrachial dystonic seizures precede LGI1 antibody limbic encephalitis. Ann Neurol 2011;69:892–900 CrossRef Medline
- Gadoth A, Pittock SJ, Dubey D, et al. Expanded phenotypes and outcomes among 256 LGI1/CASPR2-IgG-positive patients. Ann Neurol 2017;82:79–92 CrossRef Medline
- Cousyn L, Lambrecq V, Houot M, et al. Seizures in autoimmune encephalitis: specific features based on a systematic comparative study. *Epileptic Disord* 2021;23:879–92 CrossRef Medline
- 52. Kambadja B, Marion H, Cousyn L, et al. When should we test patients with epilepsy for autoimmune antibodies? Results from a French retrospective single center study. *J Neurol* 2022;269:3109–3118 CrossRef Medline
- Sola-Valls N, Ariño H, Escudero D, et al. Telemedicine assessment of long-term cognitive and functional status in anti-leucine-rich, glioma-inactivated 1 encephalitis. Neurol Neuroimmunol Neuroinflamm 2019;7:e652 CrossRef Medline
- 54. Huang X, Fan C, Gao L, et al. Clinical features, immunotherapy, and outcomes of anti-leucine-rich glioma-inactivated-1 encephalitis. J Neuropsychiatry Clin Neurosci 2022;34:141–48 CrossRef Medline
- 55. Teng Y, Li T, Yang Z, et al. Clinical features and therapeutic effects of anti-leucine-rich glioma inactivated 1 encephalitis: a systematic review. Front Neurol 2021;12:791014 CrossRef Medline
- 56. Zhang T, Duan Y, Ye J, et al. Brain MRI characteristics of patients with anti-N-methyl-D-aspartate receptor encephalitis and their associations with 2-year clinical outcome. AJNR Am J Neuroradiol 2018;39:824–29 CrossRef Medline
- 57. Joubert B, Saint-Martin M, Noraz N, et al. Characterization of a subtype of autoimmune encephalitis with anti-contactin-associated protein-like 2 antibodies in the cerebrospinal fluid, prominent limbic symptoms, and seizures. *JAMA Neurol* 2016;73:1115–24 CrossRef Medline
- Moura J, Samões R, Cardoso M, et al. Distinct phenotypes in a cohort of anti-CASPR2 associated neurological syndromes. *Clin Neurol Neurosurg* 2023;234:107994 CrossRef Medline
- 59. Zhu F, Shan W, Lv R, et al. Clinical characteristics of anti-GABA-B receptor encephalitis. *Front Neurol* 2020;11:403 CrossRef Medline
- 60. Höftberger R, Titulaer MJ, Sabater L, et al. Encephalitis and GABAB receptor antibodies: novel findings in a new case series of 20 patients. *Neurology* 2013;81:1500–06 CrossRef Medline

- 61. Gresa-Arribas N, Planagumà J, Petit-Pedrol M, et al. Human neurexin-3α antibodies associate with encephalitis and alter synapse development. *Neurology* 2016;86:2235–42 CrossRef Medline
- 62. Höftberger R, van Sonderen A, Leypoldt F, et al. Encephalitis and AMPA receptor antibodies: novel findings in a case series of 22 patients. *Neurology* 2015;84:2403–12 CrossRef Medline
- 63. Joubert B, Kerschen P, Zekeridou A, et al. Clinical spectrum of encephalitis associated with antibodies against the α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor: case series and review of the literature. JAMA Neurol 2015;72:1163–69 CrossRef Medline
- 64. Spatola M, Sabater L, Planagumà J, et al. Encephalitis with mGluR5 antibodies: symptoms and antibody effects. *Neurology* 2018;90: e1964–72 CrossRef Medline
- 65. Carvajal-González A, Leite MI, Waters P, et al. Glycine receptor antibodies in PERM and related syndromes: characteristics, clinical features and outcomes. *Brain* 2014;137:2178–92 CrossRef Medline
- 66. Swayne A, Tjoa L, Broadley S, et al. Antiglycine receptor antibody related disease: a case series and literature review. Eur J Neurol 2018;25:1290–98 CrossRef Medline
- Muñiz-Castrillo S, Hedou JJ, Ambati A, et al. Distinctive clinical presentation and pathogenic specificities of anti-AK5 encephalitis. *Brain* 2021;144:2709–21 CrossRef Medline
- Xiao J, Fu PC, Li ZJ. Clinical and imaging analysis to evaluate the response of patients with anti-DPPX encephalitis to immunotherapy. *BMC Neurol* 2022;22:129 CrossRef Medline
- Fredriksen JR, Carr CM, Koeller KK, et al. MRI findings in glutamic acid decarboxylase associated autoimmune epilepsy. *Neuroradiology* 2018;60:239–45 CrossRef Medline
- 70. Sun Y, Qin X, Huang D, et al. Anti-amphiphysin encephalitis: expanding the clinical spectrum. *Front Immunol* 2023;14:1084883 CrossRef Medline
- Dalmau J, Graus F, Villarejo A, et al. Clinical analysis of anti-Ma2associated encephalitis. Brain 2004;127:1831–44 CrossRef Medline
- 72. Quintas-Neves M, Teylan MA, Morais-Ribeiro R, et al. Divergent magnetic resonance imaging atrophy patterns in Alzheimer's disease and primary age-related tauopathy. *Neurobiol Aging* 2022;117:1–11 CrossRef Medline
- 73. Pinho J, Almeida FC, Araújo JM, et al. Sex-specific patterns of cerebral atrophy and enlarged perivascular spaces in patients with cerebral amyloid angiopathy and dementia. AJNR Am J Neuroradiol 2023;44:792–98 CrossRef Medline
- 74. Preiß D, Billette OV, Schneider A, et al. The atrophy pattern in Alzheimer-related PPA is more widespread than that of the frontotemporal lobar degeneration associated variants. *NeuroImage Clin* 2019;24:101994 CrossRef Medline
- Nelson PT, Dickson DW, Trojanowski JQ, et al. Limbic-predominant age-related TDP-43 encephalopathy (LATE): consensus working group report. Brain 2019;142:1503–27 CrossRef Medline
- 76. Eran A, Hodes A, Izbudak I. Bilateral temporal lobe disease: looking beyond herpes encephalitis. *Insights Imaging* 2016;7:265–74 CrossRef Medline
- 77. de Priester JA, Jansen GH, de Kruijk JR, et al. New MRI findings in Creutzfeldt-Jakob disease: high signal in the globus pallidus on T1-weighted images. Neuroradiology 1999;41:265–68 CrossRef Medline
- 78. de Oliveira AM, Paulino MV, Vieira AP, et al. Imaging patterns of toxic and metabolic brain disorders. *Radiographics* 2019;39:1672–95 CrossRef Medline
- 79. Fragoso DC, Gonçalves Filho AL, Pacheco FT, et al. Imaging of Creutzfeldt-Jakob disease: imaging patterns and their differential diagnosis. *Radiographics* 2017;37:234–57 CrossRef Medline
- 80. Vakrakou A, Constantinides VC, Velonakis G, et al. Paraneoplastic basal ganglia encephalitis associated with anti-CV2/CRMP-5 and anti-Yo antibodies in a patient with non-small-cell lung cancer. *Neurol Sci* 2020;41:2649–51 CrossRef Medline

- Hegde AN, Mohan S, Lath N, et al. Differential diagnosis for bilateral abnormalities of the basal ganglia and thalamus. *Radiographics* 2011;31:5–30 CrossRef Medline
- Miller TD, Chong TT, Aimola Davies AM, et al. Focal CA3 hippocampal subfield atrophy following LG11 VGKC-complex antibody limbic encephalitis. *Brain* 2017;140:1212–19 CrossRef Medline
- Petit-Pedrol M, Sell J, Planagumà J, et al. LGI1 antibodies alter Kv1.1 and AMPA receptors changing synaptic excitability, plasticity and memory. *Brain* 2018;141:3144–59 CrossRef Medline
- Ramberger M, Berretta A, Tan JM, et al. Distinctive binding properties of human monoclonal LGI1 autoantibodies determine pathogenic mechanisms. *Brain* 2020;143:1731–45 CrossRef Medline
- 85. Kornau HC, Kreye J, Stumpf A, et al. Human cerebrospinal fluid monoclonal lgi1 autoantibodies increase neuronal excitability. *Ann Neurol* 2020;87:405–18 CrossRef Medline
- 86. Sadaghiani MS, Roman S, Diaz-Arias LA, et al. Comparison of quantitative FDG-PET and MRI in anti-LGI1 autoimmune encephalitis. *Neuroradiology* 2023;65:1225–38 CrossRef Medline