

Supplementary Table 1. Clinical characteristics of included patients with idiopathic intracranial hypertension (IIH).

Clinical Variable	Number (%)
<i>Patient Demographics</i>	
Total IIH patients	99 (100)
Female	88 (89)
Age at diagnosis	29 (SD: 8)
mean BMI	40.7 (SD: 8.3)
White	68 (68.7)
Black	23 (23.2)
Other Ethnicity	8 (8.1)
<i>Clinical Features</i>	
Headache	90 (91)
Tinnitus	43 (43)
Visual Disturbance	69 (69.7)
Papilledema	93 (94)
Mean opening pressure	37.6 (SD: 9.4)
<i>MR Venogram</i>	91 (92)
Venous Sinus Stenosis	82 (90)
<i>Frisen Papilledema Grade</i>	
Not documented	18 (18.2)
0	6 (6.1)
1	19 (19.2)
2	27 (27.3)
3	19 (19.2)
4	8 (8.1)
5	2 (2)
<i>Past Medical History</i>	
Hypertension	21 (21.2)
Hyperlipidemia	4 (4)
Diabetes Mellitus	6 (6.1)
Thyroid Disorder	3 (3)
<i>Relevant Medications</i>	
Diuretics	10 (10.1)
Steroids	1 (0.01)
Contraceptives	6 (6.1)
Antibiotics	4 (4)
Lithium	0 (0)
Vitamin A	0 (0)
Anticoagulation	0 (0)