

On-line Table: Baseline demographics and clinical characteristics among patients who underwent endovascular treatment categorized according to procedure times

	Patients with Procedure Time ≤30 Minutes	Patients with Procedure Time >30 Minutes	P Value
Total	65	144	
Women	24 (36.9)	76 (52.7)	.0337
Age (yr) (mean)	63.6 ± 15.26	66.2 ± 16.09	.6363
Time interval between onset and microcatheter (mean)	311.8 ± 104.8	294.5 ± 113.7	.471
Time interval between onset and recanalization (mean)	330.4 ± 102.1	358.8 ± 115.4	.3179
Time interval between microcatheter to recanalization (mean)	18.8 ± 7.02	64.1 ± 31.05	<.0001
Admission NIHSS score strata			
NIHSS score 0–9	17 (26.56)	23 (15.97)	.1932
NIHSS score 10–19	33 (51.56)	82 (56.94)	
NIHSS score 20–42	14 (21.88)	39 (27.08)	
Treated artery			
Anterior cerebral artery	2 (3.1)	6 (4.17)	.7751
Internal carotid artery	9 (13.8)	31 (21.5)	
Middle cerebral artery	49 (75.3)	100 (69.5)	
Posterior cerebral artery	2 (3.1)	2 (1.4)	
Basilar artery	3 (4.5)	5 (3.5)	
Type of treatment			
Intra-arterial thrombolytic administration	41 (63.08)	114 (79.17)	.0139
Mechanical thrombectomy	17 (26.15)	60 (41.67)	.0314
Stent placement	10 (15.38)	22 (15.28)	.9842
Intra-arterial thrombolytic administration and mechanical thrombectomy	10 (15.4)	47 (32.6)	.0095
Intra-arterial thrombolytic administration plus stent placement	1 (1.5)	8 (5.6)	.1854
Angiographic recanalization	60 (92.31)	116 (80.56)	.031
Intravenous thrombolytic administered	23 (35.94)	64 (44.44)	.251
ICH symptomatic	2 (3.08)	11 (7.64)	.2062
ICH asymptomatic	3 (4.62)	12 (8.33)	.3351
Unfavorable outcome at discharge (mRS 3–6)	34 (52.31)	104 (72.22)	.0049