

On-line Table: Emergency ICA stenting in concert with MT: review of the literature (only studies with $n > 20$)

	This Study, 2015 ($n = 170$)	Lockau et al, 2015 ⁵ ($n = 37$)	Heck and Brown, 2015 ⁶ ($n = 23$)	Maurer et al, 2014 ⁷ ($n = 43$)	Cohen et al, 2015 ¹⁰ ($n = 24$)	Stampfl et al, 2014 ⁸ ($n = 24$)	Puri et al, 2015 ²² ($n = 28$)	Malik et al, 2011 ⁹ ($n = 77$)
sICH (n/N) (%)	15/170 (9%)	4/37 (11%)	5/23 (22%) ^a	NR	0/24 (0%)	4/24 (17%)	2/28 (7%)	8/77 (10%) ^e
In-hospital mortality (n/N) (%)	32/170 (19%)	7/37 (19%)	9/23 (39%) ^b	9/43 (21%)	2/24 (8%) ^b	4/24 (17%) ^b	NA	18/77 (23%)
TICI $\geq 2b$, (n/N) (%)	130/170 (77%)	27/37 (73%)	17/23 (74%)	33/43 (77%)	19/24 (79%)	15/24 (63%)	20/28 (71%)	58/77 (75%) ^f
FU mRS ≤ 2 (n/N) (%)	62/170 (36%)	17/37 (46%)	12/23 (52%)	14/43 (33%) ^c	13/17 (76%) ^d	7/24 (29%)	13/23 (57%)	32/77 (42%)

Note:—NA indicates not applicable; NR, not reported; FU, follow-up.

^a Safe Implementation of Thrombolysis in Stroke-Monitoring Criteria.

^b 90 days.

^c At discharge.

^d Of the living patients at 3 months.

^e Any PH.

^f $\geq$ TIMI 2.