

On-line Table: Clinical features and MRA findings of SEAVFs with intradural venous reflux

Clinical Features										MRA Findings		
Numbness and Tingling in Bilateral Lower Extremities												
No.	Age (y), Sex	Duration of Symptoms (mo)	Progressive Paraparesis	Urinary Bladder Dysfunction	Low Backache	Arterialized Epidural Venous Pouch (Epidural Fistula)	Additional Arterialized Epidural Veins	Point of Reflux into Radicular Vein	MRA and DSA Concordant?			
1	58, M	12	+	—	—	L3, left anterolateral	+	Superior aspect of pouch	Yes			
2	67, M	0.5	+	+	+	L4, left anterolateral	+	Epidural venous system at higher (L1–L2) level than pouch	Yes			
3	58, M	6	+	+	—	L3, left anterolateral	—	Superior aspect of pouch	Yes			
4	75, M	1.5	+	—	—	L2, left anterolateral	—	Inferior aspect of pouch	Yes			
5	54, M	2	+	—	+	L3, right anterolateral	+	Superior aspect of pouch	Feeders missed on MRA (bilateral L3, right L2, right lateral sacral feeders on DSA)			
6	62, M	4	+	+	+	L3, right anterolateral	+	Superior aspect of pouch	Contralateral feeder missed on MRA (bilateral L3 feeders on DSA)			
7	58, M	5	+	+	—	C4, left anterolateral	—	Inferior aspect of pouch	Yes			

Note:— indicates absent; +, present.